

IMPACT OF AMMONIUM NITRATE ON WOOD FROG (*RANA SYLVATICA*) TADPOLES: EFFECTS ON SURVIVORSHIP AND BEHAVIOR

AMBER A. BURGETT¹, CHRISTIAN D. WRIGHT, GEOFFREY R. SMITH², DORAN T. FORTUNE, AND
SAMUEL L. JOHNSON

Department of Biology, Denison University, Granville, Ohio 43023, USA

¹Present Address: Department of Biology, Washington University, St. Louis, Missouri 63130, USA

²Author for Correspondence: smithg@denison.edu

Abstract.—Nitrogenous based fertilizers, such as ammonium nitrate, are commonly used in agriculture, entering aquatic ecosystems through runoff. Ammonium nitrate has been shown to affect the survivorship and behavior of anurans. We conducted an experiment to examine the potential toxic effects of ammonium nitrate on Wood Frog (*Rana sylvatica*) tadpoles. We also examined whether ammonium nitrate might interact with predator cues to affect tadpole behavior. Ammonium nitrate decreased survivorship of Wood Frog tadpoles ($\geq 50 \text{ mg L}^{-1} \text{ NH}_4\text{NO}_3$). Activity level of Wood Frog tadpoles decreased when exposed to ammonium nitrate, as well as in the presence of predator (Mosquitofish, *Gambusia affinis*) chemical cues. Our results suggest that ammonium nitrate can have significant effects on anuran larvae, both through direct toxicological effects on survivorship, but also through behavioral effects.

Key Words.—activity; ammonium nitrate; behavior; *Gambusia affinis*; Mosquitofish; *Rana sylvatica*; toxicology; Wood Frog

INTRODUCTION

Nitrogenous based fertilizers are commonly used in agriculture, entering ponds and other bodies of water through runoff associated with agriculture, industrial and human wastes, livestock waste, lawn fertilizers (Fenn et al. 2003; Holland et al. 2005). The amount of these fertilizers entering freshwater ecosystems is likely to remain the same, or even increase in the future (e.g., Tilman et al. 2001).

Several studies have examined the direct effects of nitrogen fertilizers on larval amphibians. Nitrogenous compounds are known to be lethal to some species of amphibian larvae at or near ecologically relevant concentrations (e.g., Marco et al. 1999; Ortiz et al. 2004; Smith et al. 2005; see also an earlier review by Rouse et al. 1999), but not always (e.g., Allran and Karasov 2000; Hatch and Blaustein 2000; Vaala et al. 2004).

Fewer studies have examined the sublethal effects of nitrogenous compounds on larval amphibians. Such sublethal effects can include decreased growth (e.g., Allran and Karasov 2000; Johansson et al. 2001; Smith et al. 2005), changes in the timing of metamorphosis (e.g., Edwards et al. 2006), and changes in behavior (e.g., Marco and Blaustein 1999; Marco et al. 1999; Hatch and Blaustein 2000). Sublethal effects have the potential to affect interactions between larval amphibians and their competitors and predators. In many instances, sublethal effects of a variety of pollutants have been shown to affect the community structure of ponds by altering the antipredator response of amphibian larvae (e.g., Bridges 1999; Lefcort et al. 1999; Rohr and Crumrine 2005) but they may also affect competition (e.g., Smith et al. 2006). Thus, pollutants have the potential to alter the interactions between amphibians and their predators.

We studied the effects of ammonium nitrate on the survivorship and behavior (activity level in the presence and absence of cues from a potential predator, the Mosquitofish *Gambusia affinis*) of Wood Frog (*Rana sylvatica*) tadpoles.

Nitrogenous fertilizer run-off into surface waters frequently peaks with early spring rains or snow melt (see Rouse et al. 1999; see also discussions in Hecnar 1995; Ortiz-Santaliestra et al. 2006); thus, early spring breeders, such as Wood Frogs, might be particularly affected by nitrogen-based fertilizers, such as ammonium nitrate. Previous studies have shown that Wood Frog tadpoles are affected, either lethally or sublethally, by chronic exposure to ecologically relevant concentrations of nitrite (Griffis-Kyle 2005, 2007), but not to acute exposure to $\leq 20 \text{ mg L}^{-1}$ of nitrate or $\leq 6 \text{ mg L}^{-1}$ of nitrite (Smith, in press). We also examined the influence of tadpole age/size on the effects of ammonium nitrate.

MATERIALS AND METHODS

Several (> 4) wood frog egg masses were collected from a pond on the Denison University Biological Reserve, Licking Co., Ohio and brought into the laboratory to hatch and develop for approximately one week prior to our first experiment. Plastic containers, approximately $19 \times 14 \times 5 \text{ cm}$, containing 700 ml of solution, were used for this experiment. The tadpoles were exposed to one of four nominal ammonium nitrate treatments (replicated 5 times): control = 0 mg l^{-1} , low = 50 mg l^{-1} ($= 11.2 \text{ mg l}^{-1} \text{ NH}_4^+$, $= 38.8 \text{ mg l}^{-1} \text{ NO}_3^-$), medium = 100 mg l^{-1} ($= 22.5 \text{ mg l}^{-1} \text{ NH}_4^+$, $= 77.5 \text{ mg l}^{-1} \text{ NO}_3^-$), and high = 200 mg l^{-1} ($= 45.0 \text{ mg l}^{-1} \text{ NH}_4^+$, $= 155.0 \text{ mg l}^{-1} \text{ NO}_3^-$) made using ammonium nitrate crystals (CAS#61184-52-2; Fisher Scientific, Rochester, NY) and aged tapwater (water chemistry measured on aged tapwater used: Salinity = 0.3 ppt; pH = 8.5; dissolved oxygen = 9.40 mg l^{-1} ; Nitrate-N $< 1 \text{ ppm}$; Ammonia-N $\leq 0.1 \text{ ppm}$; Phosphate-P $\leq 1 \text{ ppm}$; Hardness = 176 ppm). These concentrations are within observed concentrations of nitrate or ammonium nitrate found in nature ($10\text{--}250 \text{ mg l}^{-1}$; Xu and Oldham 1997; Rouse et al. 1999; Ortiz et al. 2004) and tested on other species of anuran tadpoles (see Table 1). Six tadpoles were placed in each replicate container creating a density of $8.6 \text{ tadpoles l}^{-1}$, which is at




FIGURE 1. Effect of ammonium nitrate concentrations and tadpole age (early vs. late – see text) on the survivorship of Wood Frog (*Rana sylvatica*) tadpoles for one week. Means are given ± 1 SE $N = 5$ replicates for all means.

the lower end of the range of densities used in other experiments on ammonium nitrate (see Table 1).

Tadpoles were exposed to solutions for one week. All tadpoles were fed the same amount of ground Purina Rabbit Chow on the first day of exposure and on the fourth day of exposure. Excess food and feces were removed before the food was added on the fourth day. After the week of exposure, survivorship was determined using the number of tadpoles still alive in each container. Behavioral observations began after the survivorship data were collected. We recorded the number of movements of two focal tadpoles per container for ten minutes. Containers were divided into four quadrants, and we considered a tadpole to have moved each time it crossed a line into a different quadrant. After these initial observations were made, we added 50 ml of water from a holding tank (water volume = 5 l) occupied by 20 *G. affinis* for at least 1 week. Tadpoles were then given ten minutes to re-acclimate, and then observations on activity level were repeated for two focal tadpoles in each container as outlined above. We thus had pre- and post-cue activity level observations for each container (see Kruuk and Gilchrist 1997; Marquis et al. 2004 for similar methods).

A second set of experiments and behavioral observations (“late experiment”) were conducted two weeks after the first set of experiments to allow for growth and development of the tadpoles. The two sets of experiments used the same methods. Tadpoles used in the second experiment were not exposed to ammonium nitrate until the start of the second experiment. Mean mass of 10 haphazardly selected tadpoles at the start of the early experiment was 0.080 ± 0.0005 g. Mean mass of 10 haphazardly selected tadpoles at the start of the late experiment was 0.094 ± 0.009 g. In both experiments tadpoles were Gosner stage 26 (Gosner 1960).

To examine the effects of tadpole age (early vs. late) and the nominal concentration of ammonium nitrate (control, low, medium, or high) on tadpole survivorship, we used a two-way ANOVA ($\alpha = 0.05$) on the arcsin-square root transformed survivorships (i.e., proportion of tadpoles surviving for each replicate). We used repeated-measure ANOVAs ($\alpha = 0.05$) to examine the effects of predator cue (as a repeated measure) and ammonium nitrate on the activity levels of the tadpoles. Preliminary analyses of activity using the number of surviving tadpoles as a covariate found no significant effects of the covariate and no significant interactions of the covariate and ammonium nitrate concentration. Due to the complete mortality

of tadpoles at the high concentration in the early experiment, we used separate analyses of the activity levels for the two experiments.

RESULTS

Survivorship.—Survivorship of tadpoles significantly decreased as the concentration of ammonium nitrate increased (Fig. 1; $F_{3,32} = 23.7$, $P < 0.0001$; all pairwise Fisher’s PLSD tests among ammonium nitrate groups were significant at $P \leq 0.003$, except for the low vs. medium concentration comparison, which was not significant). Tadpole age had no effect on survivorship (Fig. 1; $F_{1,32} = 0.56$, $P = 0.46$). However, the interaction of treatment and age did show a significant effect on tadpole survivorship. Mortality at low and medium concentrations of ammonium nitrate was slightly higher in the late experiment than in the early experiment; whereas, at the high concentration, mortality was complete in the early experiment, but not in the late experiment (Fig. 1, $F_{3,32} = 5.99$, $P = 0.0023$).

Activity Level.—For the early experiment, activity was, in general, lower at low and medium concentrations of ammonium nitrate than in the control (Fig. 2A; $F_{2,12} = 8.98$, $P = 0.0041$; Fisher’s PLSD: control vs. low, $P = 0.0012$; control vs. medium, $P = 0.026$). Activity was also significantly lower in the presence of the potential predator cue (Fig. 2A; $F_{1,12} = 7.46$, $P = 0.018$). There was no significant interaction between ammonium nitrate concentration and the presence of the potential predator cue ($F_{2,12} = 1.96$, $P = 0.18$).

For the late experiment, activity was not significantly affected by ammonium nitrate concentration, although activity levels did tend to be lower in the presence of ammonium nitrate (Figure 2B; $F_{3,14} = 1.43$, $P = 0.28$). Activity tended to be lower in the presence of the potential predator cue, but the difference was not




FIGURE 2. Effects of ammonium nitrate concentrations and the presence of a predator (*Gambusia affinis*) cue on the activity of Wood Frog (*Rana sylvatica*) tadpoles in the (A) early experiment (total mortality at the high concentration precluded measuring of activity), and (B) late experiment. Means are given ± 1 SE.

TABLE 1.— A review of the susceptibility of anuran tadpoles for exposure to ammonium nitrate. Values given are mg L⁻¹ of ammonium nitrate unless otherwise noted.

Species	Duration	Lethal Effects (> 50% mortality)	Not lethal (< 25% mortality for all treatments)	Tadpole Density [tadpolesl ⁻¹]	Source
		[mg l ⁻¹ NH ₄ NO ₃]	[mg l ⁻¹ NH ₄ NO ₃]		
<i>Bufo americanus</i>	96 h	80-228		10	Hecnar (1995)
<i>B. bufo</i>	15 d	1143		12-20	Ortiz et al. (2004)
<i>B. bufo</i>	96 h-7 d	≈ 2193		12.5	Xu and Oldham (1997)
<i>B. calanita</i>	8-12 d	648		?	Ortiz-Santaliestra et al. (2006)
<i>B. calanita</i>	15 d		Up to 1143	12-20	Ortiz et al. (2004)
<i>Crinia signifera</i>	21 d		Up to 19.4	19.8	Hamer et al. (2004)
<i>Discoglossus galganoi</i>	15 d	259		?	Ortiz-Santaliestra et al. (2006)
<i>D. galganoi</i>	15 d	1143		12-20	Ortiz et al. (2004)
<i>Hyla arborea</i>	15 d	286		12-20	Ortiz et al. (2004)
<i>Limnodynastes peronii</i>	91 d		Up to 19.4	39.6	Hamer et al. (2004)
<i>Litoria aurea</i>	150 d	12.9		9.9	Hamer et al. (2004)
<i>Osteopilus septentrionalis</i>	13 d	228.6		2	Punzo and Law (2006)
<i>Pelobates cultripes</i>	8-15 d	259		?	Ortiz-Santaliestra et al. (2006)
<i>P. cultripes</i>	15 d		Up to 1143	12-20	Ortiz et al. (2004)
<i>Pleurodeles waltl</i>	15 d		Up to 1143	12-20	Ortiz et al. (2004)
<i>Pseudacris regilla</i>	10 d	281-570		4 – 6	Schuytema and Nebeker (1999)
<i>P. triseriata</i>	96 h	143		10	Hecnar (1995)
<i>Rana clamitans</i>	96 h	> 286		10	Hecnar (1995)
<i>R. pipiens</i>	96 h	143		10	Hecnar (1995)
<i>R. sylvatica</i>	7 d	50-100		8.6	This Study
<i>Xenopus laevis</i>	10 d	281-570		15	Schuytema and Nebeker (1999)

statistically significant (Fig. 2B; $F_{1,14} = 3.69$, $P = 0.075$). There was no significant interaction between ammonium nitrate concentration and the predator cue ($F_{3,14} = 1.41$, $P = 0.28$).

DISCUSSION

Our study suggests that ammonium nitrate can have a significant negative effect on the survivorship of Wood Frog tadpoles, at least at the range of concentrations used in this experiment. Our findings place the susceptibility of Wood Frog tadpoles to ammonium nitrate within the range of susceptibilities to ammonium nitrate found for a wide range of other species of anurans (Table 1). Thus, while Wood Frogs may not be physiologically more prone to ecological effects of ammonium nitrate exposure, they may be at more risk than other anurans if, as previous studies have suggested (see Hecnar 1995; Rouse et al 1999; Ortiz-Santaliestra et al. 2006), fertilizer runoff and contamination of surface waters peaks in the late winter or early spring when Wood Frogs breed.

Our results suggest that the negative effects of exposure to ammonium nitrate may change with age in Wood Frog tadpoles (e.g., total mortality at high concentrations in the early experiment, but 33% survivorship during the late experiment). The age and size differences in our study were relatively small, and all the tadpoles were the same developmental stage when tested. However, Ortiz-Santaliestra et al. (2006) found that a difference in only four days was enough to change the susceptibility of the hatchlings of *Pelobates cultripes* to ammonium nitrate, suggesting that small changes in age or size may be able to affect susceptibility of anuran tadpoles to ammonium nitrate. Other studies have also found tolerance for exposure to toxic substances can change over the development or growth of tadpoles (e.g., Harris et al. 2000; Smith 2001).

Our results show that Wood Frog tadpoles tend to decrease activity when exposed to ammonium nitrate at both early and late

exposure times (albeit only significantly in the early experiment). Exposure to ammonium nitrate has been shown to reduce activity or feeding rate in a variety of other amphibian larvae (e.g., Hecnar 1995; Watt and Oldham 1995; Xu and Oldham 1997). The decrease in activity level of tadpoles or larvae in the presence of ammonium nitrate may actually decrease predation risk since activity in tadpoles frequently increases predation (see Lefcort 1996; Anholt et al. 2005). However, reduced activity could have negative effects since foraging, and consequently growth, are often functions of activity levels (Skelly and Werner 1990; Relyea and Werner 1999). Reduced activity may also affect the competitive abilities of tadpoles because activity level has sometimes been correlated with competitive ability in tadpoles (e.g., Morin and Johnson 1988; Werner 1992; Semlitsch 1993; Dayton and Fitzgerald 2001), but not always (e.g., Smith et al. 2004). Thus the consequences of these sublethal, behavioral effects of ammonium nitrate on Wood Frog tadpoles may depend on the ecological context of the exposure (e.g., whether predators or competitors are present or not).

Wood Frog tadpoles have been shown to decrease activity in the presence of fish (*Umbra limi*, Relyea 2000; *Lepomis gibbosus*, Richardson 2001; *L. macrochirus*, A.R. Awan and G.R. Smith In Press). In general, Wood Frog tadpoles in our experiment showed a decrease in activity level when exposed to water containing chemical cues from a potential predator, *G. affinis*, but this reduction in activity was only statistically significant in the early experiment. Size does not appear to affect the ability of Mosquitofish to prey upon tadpoles (Webb and Joss 1997); therefore, all tadpoles in this experiment would likely be susceptible to predation by the Mosquitofish. Lawler et al. (1994) found that younger tadpoles (Gosner Stage 26) of *Rana aurora draytoni* reduced activity levels in the presence of Mosquitofish, but older tadpoles (Gosner Stage 33-36) showed no reduction in activity. Other species of tadpoles show no response to the presence of Mosquitofish cues (Hamer et al. 2002; G.R.

Smith et al. unpubl. data). There was no significant interaction between predator cue presence and exposure to ammonium nitrate. Thus, there was no evidence that exposure to ammonium nitrate modified the response of tadpoles to the cues of Mosquitofish.

Acknowledgments.—Egg masses were collected under a permit from the Ohio Department of Natural Resources, and the experiments conducted with approval by the Denison University IACUC (#05-005; #06-007).

LITERATURE CITED

- Allran, J.W., and W.H. Karasov. 2000. Effects of atrazine and nitrate on Northern Leopard Frog (*Rana pipiens*) larvae exposed in the laboratory from posthatch through metamorphosis. *Environmental Toxicology and Chemistry* 19:2850-2855.
- Anholt, B.R., S. Negovetic, C. Ranta, and C. Som. 2005. Predator complement determines the relative success of tadpoles of the *Rana esculenta* complex. *Evolutionary Ecology Research* 7: 733-741.
- Awan, A.R. and G.R. Smith. In Press. The effect of group size on the responses of Wood Frog tadpoles to fish. *American Midland Naturalist*.
- Bridges, C.M. 1999. Predator-prey interactions between two amphibian species: Effects of insecticide exposure. *Aquatic Ecology* 33:205-211.
- Dayton, G.H., and L.A. Fitzgerald. 2001. Competition, predation, and the distributions of four desert anurans. *Oecologia* 129: 430-435.
- Edwards, T.M., K.A. McCoy, T. Barbeau, M.W. McCoy, J.M. Thro, and L.J. Guillelte Jr. 2006. Environmental context determines nitrate toxicity in Southern Toad (*Bufo terrestris*) tadpoles. *Aquatic Toxicology* 78:50-58.
- Fenn, M.E., J.S. Baron, E.B. Allen, H.M. Rueth, K.R. Nydick, and L. Geiser. 2003. Ecological effects of nitrogen deposition in the western United States. *BioScience* 53:404-420.
- Gosner, K.L. 1960. A simplified table for staging anuran embryos and larvae with notes on identification. *Herpetologica* 16:183-190.
- Griffis-Kyle, K.L. 2005. Ontogenetic delays in effects of nitrite exposure on Tiger Salamanders (*Ambystoma tigrinum tigrinum*) and Wood Frogs (*Rana sylvatica*). *Environmental Toxicology and Chemistry* 24:1523-1527.
- Griffis-Kyle, K.L. 2007. Sublethal effects of nitrite on Eastern Tiger Salamander (*Ambystoma tigrinum tigrinum*) and Wood Frog (*Rana sylvatica*) embryos and larvae: implications for field populations. *Aquatic Ecology* 41:119-127.
- Hamer, A.J., S.J. Lane, and M.J. Mahony. 2002. The role of introduced Mosquitofish (*Gambusia holbrooki*) in excluding the native Green and Golden Bell Frog (*Litoria aurea*) from original habitats in south-eastern Australia. *Oecologia* 132:445-452.
- Hamer, A.J., J.A. Makings, S.J. Lane, and M.J. Mahony. 2004. Amphibian decline and fertilizers used on agricultural land in south-eastern Australia. *Agriculture, Ecosystems and Environment* 102:299-305.
- Harris, M.L., L. Chora, C.A. Bishop, and J.P. Bogart. 2000. Species- and age-related differences in susceptibility to pesticide exposure for two amphibians, *Rana pipiens*, and *Bufo americanus*. *Bulletin of Environmental Contamination and Toxicology* 64:263-270.
- Hatch, A.C., and A.R. Blaustein. 2000. Combined effects of UV-B, nitrate, and low pH reduce the survival and activity level of larval Cascades Frogs (*Rana cascadae*). *Archives of Environmental Contamination and Toxicology* 39:494-499.
- Hecnar, S.J. 1995. Acute and chronic toxicity of ammonium nitrate fertilizer to amphibians from southern Ontario. *Environmental Toxicology and Chemistry* 14:2131-2137.
- Holland, E.A., B.H. Braswell, J. Sulzman, and J.-F. Lamarque. 2005. Nitrogen deposition onto the United States and western Europe: synthesis of observations and models. *Ecological Applications* 15:38-57.
- Johansson, M., K. Räsänen, and J. Merilä. 2001. Comparison of nitrate tolerance between different populations of the Common Frog, *Rana temporaria*. *Aquatic Toxicology* 54:1-14.
- Kruuk, L.E.B., and J.S. Gilchrist. 1997. Mechanism maintaining species differentiation: predator-mediated selection in a *Bombina* hybrid zone. *Proceedings of the Royal Society of London* 264B:105-110.
- Lawler, S.P., D. Dritz, T. Strange, and M. Holyoak 1999. Effects of introduced Mosquitofish and Bullfrogs on the threatened California Red-legged Frog. *Conservation Biology* 13:613-622.
- Lefcort, H. 1996. Adaptive, chemically mediated fright response in tadpoles of the Southern Leopard Frog, *Rana utricularia*. *Copeia* 1996:455-459.
- Lefcort, H., S.M. Thomson, E.E. Cowles, H.L. Harowicz, B.M. Livaudais, W.E. Roberts, and W.F. Ettinger. 1999. Ramifications of predator avoidance: Predator and heavy metal-mediated competition between tadpoles and snails. *Ecological Applications* 9:1477-1489.
- Marco, A., and A.R. Blaustein. 1999. The effects of nitrite on behavior and metamorphosis in Cascades Frogs (*Rana cascadae*). *Environmental Toxicology and Chemistry* 18:946-949.
- Marco, A., C. Quilchano, and A.R. Blaustein. 1999. Sensitivity to nitrate and nitrite in pond-breeding amphibians from the Pacific Northwest, USA. *Environmental Toxicology and Chemistry* 18: 2836-2839.
- Marquis, O., P. Saglio, and A. Neveu. 2004. Effects of predator and conspecific chemical cues on the swimming activity of *Rana temporaria* and *Bufo bufo* tadpoles. *Archiv für Hydrobiologie* 160:153-170.
- Morin, P.J., and E.A. Johnson. 1988. Experimental studies of asymmetric competition among anurans. *Oikos* 53:398-407.
- Ortiz, M.E., A. Marco, N. Saiz, and M. Lizana. 2004. Impact of ammonium nitrate on growth and survival of six European amphibians. *Archives of Environmental Contamination and Toxicology* 47:234-239.
- Ortiz-Santaliestra, M.E., A. Marco, M.J. Fernández, and M. Lizana. 2006. Influence of developmental stage on sensitivity to ammonium nitrate of aquatic stages of amphibians. *Environmental Toxicology and Chemistry* 25:105-111.
- Punzo, F., and S. Law. 2006. Effect of nitrate-related compounds on growth, survival and hematological responses in tadpoles of the Cuban Tree Frog, *Osteopilus septentrionalis* (Boulenger). *Journal of Environmental Biology* 27:187-190.
- Relyea, R.A. 2000. Trait-mediated indirect effects in larval anurans: reversing competition with the threat of predation. *Ecology* 81:2278-2289.

- Relyea, R.A., and E.E. Werner. 1999. Quantifying the relation between predator-induced behavior and growth performance in larval anurans. *Ecology* 80:2117-2124.
- Richardson, J.M.L. 2001. A comparative study of activity levels in larval anurans and response to the presence of different predators. *Behavioral Ecology* 12:51-58.
- Rohr, J.R., and P.W. Crumrine. 2005. Effects of an herbicide and an insecticide on pond community structure and processes. *Ecological Applications* 15:1135-1147.
- Rouse, J.D., C.A. Bishop, and J. Struger. 1999. Nitrogen pollution: An assessment of its threat to amphibian survival. *Environmental Health Perspectives* 107:799-803.
- Schuytema, G.S., and A.V. Nebeker. 1999. Comparative toxicity of ammonium and nitrate compounds to Pacific Treefrog and African Clawed Frog tadpoles. *Environmental Toxicology and Chemistry* 18:2251-2257.
- Semlitsch, R.D. 1993. Asymmetric competition in mixed populations of tadpoles of the hybridogenetic *Rana esculenta* complex. *Evolution* 47:510-519.
- Skelly, D.K., and E.E. Werner. 1990. Behavioral and life-historical responses of larval American Toads to an odonate predator. *Ecology* 71:2313-2322.
- Smith, G.R. 2001. Effects of acute exposure to a commercial formulation of glyphosate on the tadpoles of two species of anurans. *Bulletin of Environmental Contamination and Toxicology* 67:483-488.
- Smith, G.R. In press. Lack of effect of nitrate, nitrite, and phosphate on Wood Frog (*Rana sylvatica*) tadpoles. *Applied Herpetology*.
- Smith, G.R., H.A. Dingfelder, and D.A. Vaala. 2004. Asymmetric competition between *Rana clamitans* and *Hyla versicolor* tadpoles. *Oikos* 105:626-632.
- Smith, G.R., K.G. Temple, H.A. Dingfelder, and D.A. Vaala. 2006. Effects of nitrate on the interactions of the tadpoles of two ranids (*Rana clamitans* and *R. catesbeiana*). *Aquatic Ecology* 40:125-130.
- Smith, G.R., K.G. Temple, D.A. Vaala, and H.A. Dingfelder. 2005. Effects of nitrate on the tadpoles of two ranids (*Rana catesbeiana* and *R. clamitans*). *Archives of Environmental Contamination and Toxicology* 49:559-562.
- Tilman, D., J. Fargione, B. Wolff, C. D'Antonio, A. Dobson, R. Howarth, D. Schindler, W.H. Schlesinger, D. Simberloff, and D. Swackhamer. 2001. Forecasting agriculturally driven global environment change. *Science* 292:281-284.
- Vaala, D.A., G.R. Smith, K.G. Temple, and H.A. Dingfelder. 2004. No effect of nitrate on Gray Treefrog (*Hyla versicolor*) tadpoles. *Applied Herpetology* 1:265-269.
- Watt, P.J., and R.S. Oldham. 1995. The effect of ammonium nitrate on the feeding and development of larvae of the Smooth Newt, *Triturus vulgaris* (L.), and on the behaviour of its food source, *Daphnia*. *Freshwater Biology* 33:319-324.
- Webb, C., and J. Joss. 1997. Does predation by the fish *Gambusia holbrooki* (Atheriniformes: Poeciliidae) contribute to declining frog populations? *Australian Zoologist* 30:316-324.
- Werner, E.E. 1992. Competitive interactions between Wood Frog and Northern Leopard Frog larvae: the influence of size and activity. *Copeia* 1992:26-35.
- Xu, Q., and R.S. Oldham. 1997. Lethal and sublethal effects of nitrogen fertilizer ammonium nitrate on Common Toad (*Bufo bufo*) tadpoles. *Archives of Environmental Contamination and Toxicology* 32:298-303.



AMBER A. BURGETT (left) is currently a Ph.D. student in the Department of Biology at Washington University. She received her B.S. from Denison University. Her current research focuses on the effects of habitat alteration on anuran metapopulations.

CHRISTIAN D. WRIGHT (right) is a teacher with Teach for America in Phoenix, Arizona. He received his B.S. from Denison University. After his stint with Teach for America, Christian plans on attending graduate school and conduct research on snake ecology.



GEOFFREY R. SMITH is an associate professor of biology at Denison University. He received his B.A. from Earlham College, and his Ph.D. from the University of Nebraska-Lincoln. His research examines the population and community ecology of reptiles and amphibians, and in particular, how human-induced changes in the environment are affecting them.



SAMUEL L. JOHNSON is currently a senior biology major at Denison University.



DORAN T. FORTUNE is currently a senior biology major at Denison University.