
A Preliminary Survey of the Amphibian Fauna in Nilgala Forest Area and its Vicinity, Monaragala District, Sri Lanka

D.M.S. SURANJAN KARUNARATHNA^{1,2}, U.T. INDIKA ABEYWARDENA^{3,4},
M.D. CHANDANA ASELA^{3,5}, AND L.D.C. BHATHIYA KEKULANDALA^{1,6}

¹IUCN – International Union for Conservation of Nature, Sri Lanka country office,
No. 53, Horton Place, Colombo 07, Sri Lanka

²e-mail: dmsameera@gmail.com

³The Young Zoologists' Association of Sri Lanka, Department of National Zoological Gardens, Dehiwala, Sri Lanka

⁴e-mail: tiranaya@gmail.com ⁵e-mail: chandana.asela@gmail.com ⁶e-mail: bkekulandala@gmail.com

ABSTRACT.—Herein, we present the findings of a preliminary study on amphibian diversity in Nilgala Forest Area of the Monaragala District, which is situated in the intermediate zone of Sri Lanka. We investigated the amphibian species diversity and various threats they face in this remote, unstudied area. We recorded 20 species of amphibians, comprising about 18.8% of the total Sri Lankan amphibians described to date. Several species were new records for the Nilgala Forest Area including *Philautus regius*, *P. fergusonianus*, and *Ichthyophis glutinosus*. There were seven (35%) endemic species and six (30%) nationally listed threatened species observed at the site. We also observed three unidentified amphibian species in Nilgala Forest Area. Nilgala Forest Area supports high amphibian diversity. However, human activities such as burning, illegal logging, chena cultivation, and road traffic threaten the ecology of this important forest.

Key Words.—amphibians; conservation; endemic; Nilgala Forest; Sri Lanka; threatened

INTRODUCTION

Sri Lanka and Western Ghats of India is a biodiversity hotspot, rich in herpetofaunal assemblages (Meegaskumbura et al. 2002; Bossuyt et al. 2004; Gunawardene et al. 2007). Favorable environmental factors such as high rainfall and humidity, and associated high density of undergrowth found in this region support a rich diversity of herpetofauna. About 200 species of reptiles (De Silva 2006; Samarawickrama et al. 2006; Manamendra-Arachchi et al. 2007; Wickramasinghe and Munindradasa 2007; Wickramasinghe et al. 2007) and 106 species of amphibians (belonging to 16 genera, 90 species) are endemic to the island. Among those genera, *Adenomus*, *Lankanectes* and *Nannophrys* are considered relict (De Silva 1996; Dutta and Manamendra-Arachchi 1996; Manamendra-Arachchi and Pethiyagoda 2005; Meegaskumbura and Manamendra-Arachchi 2005; Fernando et al. 2007; Pethiyagoda et al. 2006; Meegaskumbura et al. 2007). The poikilothermic nature of amphibian fauna restricts their distribution to areas with high rainfall, such as lowland wet-zone forest and montane forest areas (Giri and Chaturvedi, 2001).

Wet-zone forests harbor more than 60% of the indigenous amphibian fauna of Sri Lanka, and the southwest lowland forests have almost 90% of the endemic vertebrates (Senanayake et al. 1977; Wijesinghe and Dayawansa 2002; Bambaradeniya et al. 2003; IUCN Sri Lanka and Ministry of Environmental and Natural Resources 2007). There are few studies of the

amphibian diversity in the dry and intermediate zone forests (De Silva et al. 2004). Amphibians play a vital role in the functioning of an ecosystem. Their diet of insects and small animals contributes to control of pests in human habitats, including crops and homes (Knopf 1998). However, many people in this region treat frogs with almost the same revulsion as snake, despite their ecological importance (Manamendra-Arachchi 2000).

Most researchers consider amphibians indicator species of environmental change (McCallum and Trauth 2003; Mendelson et al. 2006; McCallum 2007). Nilgala Forest Area (NFA) is one of the largest and most important forest areas in Monaragala District. However, the amphibian fauna of NFA requires study (Karunaratna et al. 2006). Several preliminary amphibian surveys took place in NFA prior to this study (Hettige et al. 2000; De Silva et al. 2004). Our new inventory work attempted to document the amphibian fauna of NFA through extensive field surveys made in 2004 and 2005. The loss of natural forests over the past 100 years contributed to the extinction of 19 species of shrub-frogs, *Philautus* sp. (Manamendra-Arachchi 2005; Meegaskumbura et al. 2007). Poorly described distributions and life histories are among the biggest hurdles for conserving amphibians in this country and abroad (e.g., Tadevosyan 2006; Jackson et al. 2007; D'Cruze et al. 2007) because only a fraction of the amphibians present in other areas of the country is known to science. This kind of information is paramount to making wise conservation decisions (Bury




FIGURE 1. Map of the study area including its major and minor road networks in the Nilgala Forest Area and its vicinity, Monaragala District, Sri Lanka.

2006; McCallum and McCallum 2006). Hence, the biodiversity information we provide in this study is important for the conservation of Sri Lanka amphibians.

MATERIALS AND METHODS

Study site.—Nilgala Forest Area is a unique forest ecosystem covering 12,432 ha in the Bibile divisional secretariat division (Fig. 1). The major vegetation type

is lowland tropical dry-mixed evergreen forest (Gunatilleke and Gunatilleke 1990; Fig. 2). Commonly found trees include Aralu (*Terminalia chebula*), Bulu (*T. bellirica*), and Nelli (*Phyllanthus emblica*). NFA occurs between 7° 08' - 7°14' N and 81°16' - 81°20' E, approximately 11 km southeast from the town of Bibile. The altitude ranges from 200-700 m above sea level within the boundaries of the NFA (De Silva et al. 2004; Karunarathna and Karunarathna 2005). The general




FIGURE 2. Some of the habitats present at Nilgala Forest Area and its vicinity, Monaragala District, Sri Lanka. A) Paddy field, B) Savannah forest, C) Riverine forest, D) Rock-outcrop in Yadunhela.

climatic conditions in Nilgala are moderately cool throughout the year, with high humidity during the northeast monsoon season.

The average annual rainfall is about 1750 mm. Most of this occurs from the end of December to the middle of March. The weather gradually dries from the middle of August to early December with the warmest temperatures in August. The mean annual temperature in the NFA is 28°C (Range = 24-32°C). NFA is also an important major watershed for Gala Oya and Panmedilla Oya throughout the year (Karunaratna and Karunaratna 2005). There are several mountain peaks within the NFA with Yakun Hela being the highest peak (>700 m). We sampled 11 habitat types during this survey (Table 1).

Sampling methods.—We implemented this study from 24 March 2004 to 23 April 2005. We spent 56 days (10 hrs/day) in the field encompassing both the wet and dry seasons. We carried out general area surveys in the different habitat types within the NFA. These took place during the day and night. We searched a variety of habitats (e.g., water bodies, under rocks, logs and decaying vegetation, trees and bushes up to 5 m). We examined species carefully, recorded their presence, and




FIGURE 3. Species composition of amphibian families in NFA. (BD- Bufonidae / MD- Microhylidae / RD- Ranidae / ID- Ichthyophiidae).

then released them back where they were captured. When needed, we used previously published keys to identify species (Dutta and Manamendra-Arachchi 1996; Manamendra-Arachchi and Pethiyagoda 2005; and




FIGURE 4. Species richness of Amphibians in various habitat types in NFA. (HG- Home Garden / SP- Small Pond / TK- Tank / ST- Stream / RD- Road Side / SB- Shrub & Bush / PF- Paddy field / RF- Riverine forest / SF- Savannah forest / RA- Rockout area / CH- Chena)

TABLE 1. Description of major habitat types in Nilgala Forest Area and its Vicinity, Monaragala District, Sri Lanka .

Habitat Type	Description of Habitat Types
1. Chena	Trees belong to family Rutaceae are dominant and grow up to 5 m; scattered bushes present; main cultivation is maize, banana, and finger millet. Leaf litter is very low.
2. Home Garden	Mixed cropping with woody plants like a <i>Mangifera indica</i> , <i>Chloroxylon swietenia</i> , <i>Schleichera oleosa</i> , <i>Tamarindus indica</i> ; trees grows up to 15 m; shade is about 50%; leaf litter content is high and wet.
3. Paddy field	Paddy fields are moderate in extent (about 1 acre); wallowing sites are frequent along the fields; field bunds are narrow.
4. Riverine forest	Shade 80% with large tall trees growing up to 20 m; <i>Mangifera ceylanica</i> , <i>Maduca longifolia</i> , <i>Terminalia chebula</i> <i>Diospyros ebenum</i> and <i>Diospyros malabarica</i> are the dominant species; thick wet leaf litter layer available; shady forest; decaying logs are common.
5. Road Side	Generally consist of small bushes growing up to 2 m. Species such as <i>Maduca longifolia</i> , <i>Terminalia bellirica</i> and <i>Mangifera zeylanica</i> can also be found in several areas.
6. Rock-outcrop	Large rock boulders and grassy areas with seasonally moist cascade habitats. Shade 20% with tall trees. <i>Ficus mollis</i> , <i>Diospyros ebenum</i> and <i>Terminalia bellirica</i>
7. Savannah forest	Only <i>Terminalia chebula</i> , <i>Terminalia bellirica</i> and <i>Phyllanthus emblica</i> forest.
8. Shrub / Bush area	1-2 m tall randomly distributed on open soil.
9. Small Pond	Seasonally flooded, mud pond, gem pits, agricultural wells, drinking wells, mud pits, clay pits
10. Stream	Perennial flowing water bodies; 1m to 10m wide; visibility high; turbidity low.
11. Tank	Open water bodies; covered by macrophytes (25%)

Manamendra-Arachchi and Pethiyagoda 2006). Furthermore, we recorded basic environmental data at each location where we caught specimens.

RESULTS AND DISCUSSION

We recorded 20 amphibians from 14 genera (Table 2) belonging to four families residing at NFA. There were seven endemic and six nationally listed threatened species among them (IUCN Sri Lanka 2000; IUCN Sri Lanka and MENR 2007). The most abundant amphibians were ranids (*Fejervarya limnocharis*, *Philautus regius*, *Euphlyctis cyanophlyctis*, *E. hexodactyla*), while caecilians were the least encountered (Fig. 3). There were 10 terrestrial, four arboreal, four aquatic, and two fossorial species. Several captures were species records for NFA (*Philautus regius*, *P. fergusonianus*, and *Ichthyophis glutinosus*). We also recorded three species of *Philautus* (shrub-frog sp.) that are probably new to science.

Fejervarya limnocharis (12.73%) and *Philautus regius* (10.11%) were the most abundant species at NFA. The abundance of aquatic and semi aquatic habitats (small ponds, streams, and paddy fields) present at NFA is likely critical for supporting the large numbers of *Fejervarya limnocharis*; whereas, the large component of shrub vegetation here helps maintain the abundance of *Philautus regius*. It is interesting that no previously published studies recorded *P. regius* in NFA. This species is rare and or data deficient in the dry and intermediate zones (Manamendra-Arachchi and Pethiyagoda 2005).

The highest species richness occurred in home

gardens (20.78%) followed by roadside habitats (19.48%) and riverine forests (12.99%); whereas, rock outcrop habitats had the lowest species richness (2.60%) (Fig. 4). Several factors may cause the high species richness in the home gardens, including the large amount of leaf litter, shade, and the abundant of prey items (e.g., insects) attracted to artificial lighting surrounding these human-use areas. However, we sampled longer in home gardens than the in other habitats. Consequently, this may reflect a simple sampling bias in our study. Nevertheless, home gardens such as these occur in the vicinity of NFA and provide important refugia for amphibians. It appears that a higher species richness of amphibians occur in disturbed habitats such as home gardens, chena, and roadside shrub. This may be due to edge effect (Murcia 1995).

The species richness is also high along the road side (on Bibila–Ampara main road). Sampling bias may contribute to these findings as well. The lowest species richness (2.60%) occurred in the rock-outcrop habitats. Outcrops are xeric due to direct sun exposure much of the year, and are not highly suitable for amphibians (Karunaratna et al. 2006; Pough et al. 2004). Only *Nannophrys naeyakai* and *Bufo melanostictus* used rocky outcrops.

Conservation implications.—We anecdotally



FIGURE 5. A fire scorched area arising from slash-and-burn agricultural practices in Nilgala Forest Area, Monaragala District, Sri Lanka.

observed several potential threats to the amphibians occurring in the NFA. These include forest fires, illegal logging, extensive use of agrichemicals, forest clearing for chena cultivation, and roads. Local farmers use slash-and-burn (Fig. 5), cropping technologies, and they hunt throughout the area (Karunaratna et al. 2005). Fires are very frequent in the months of August and September and probably destroy the habitats suitable for amphibians. Illegal logging takes place around and within the NFA, and this seriously degrades the quality of the forests. Deforestation is known to reduce soil

TABLE 2. List of amphibian species recoded from the Nilgala Forest Area and its vicinity, Monaragala District, Sri Lanka. (Abbreviations: **CR** – Critically Endangered, **DD** – Data Deficient, **TR** – Threatened Species, * – Endemic species, ** – Endemic Genus (Source: IUCN Sri Lanka, 2000; IUCN Sri Lanka and Ministry of Environmental and Natural Resources, 2007).

Scientific Names and Family	Common Name	IUCN 2000	IUCN & MENR 2007	Total Individuals	Proportion Represented
Bufonidae					
1 <i>Bufo cf. atukoralei</i>	Atukorale's Toad *	TR		3	1.12
2 <i>Bufo melanostictus</i>	Common House Toad			17	6.37
Microhylidae					
3 <i>Kaloula taprobanica</i>	Common Bull Frog			8	3
4 <i>Microhyla ornata</i>	Ornate Narrow Mouth Frog			11	4.12
5 <i>Microhyla rubra</i>	Red Narrow Mouth Frog			13	4.87
6 <i>Ramanella variegata</i>	White Bellied Pug Snouted Frog			3	1.12
7 <i>Uperodon systoma</i>	Balloon frog			16	5.99
Ranidae					
8 <i>Hoplobatrachus crassus</i>	Jerdon's Bull Frog			19	7.12
9 <i>Fejervarya limnocharis</i>	Common Paddy Field Frog			34	12.73
10 <i>Euphlyctis cyanophlyctis</i>	Skipper Frog			24	8.99
11 <i>Euphlyctis hexodactyla</i>	Six Toed Green Frog			24	8.99
12 <i>Nannophrys naeyakai</i>	Tribble Rock Frog **		CR	14	5.24
13 <i>Rana gracilis</i>	Sri Lanka Wood Frog *	TR		4	1.5
14 <i>Sphaerotheca breviceps</i>	Banded sand frog			7	2.62
15 <i>Sphaerotheca rolendae</i>	Marbled Sand Frog			9	3.37
16 <i>Philautus fergusonianus</i>	Ferguson's Shrub Frog *	TR		3	1.12
17 <i>Philautus regius</i>	Polonnaruwa Shrub Frog *		DD	27	10.11
18 <i>Polypedates cruciger</i>	Common Hourglass Tree Frog *	TR		12	4.49
19 <i>Polypedates maculatus</i>	Chunam Tree Frog			18	6.74
Ichthyophiidae					
20 <i>Ichthyophis glutinosus</i>	Yellow Banded Caecilian *	TR		1	0.37



FIGURE 6. A road-killed *Hoplobatrachus crassus* (Jerdon's Bullfrog) observed at Nilgala Forest Area, Monaragala District, Sri Lanka.

moisture, a critical resource for maintenance of amphibian populations (Camargo and Kapos 1995).

Local paddy and chena cultivation in the vicinity of the NFA leads to agrichemical use. Such chemicals could harm many animals, including amphibians (Burgett et al. 2007; Kapfer et al. 2007; Gray et al. 2007). Another significant threat to the amphibian fauna of the area is road mortality (Fig. 6). Bibila–Ampara road cuts across the NFA, and we observed many road mortalities. We observed especially high numbers of dead amphibians after a rain. Certainly, direct and indirect effects observed in previous studies (Karraker 2007; Glista et al. 2008) could significantly impact NFA amphibians.

Our preliminary study of the amphibian fauna of the Nilgala Forest and its vicinity demonstrates that NFA is an important natural area in terms of amphibian diversity. The NFA may be an important refuge for threatened amphibians residing in the intermediate zone of the Uva province (Karunarathna et al. 2006). More detailed studies are needed to document the poorly understood amphibian community at this site, and to better understand how potentially hazardous human activities may influence this important ecological resource so that conservation officials can select appropriate mitigation strategies to conserve Nilgala Forest and its surrounding habitats.

Acknowledgements.—The authors wish to thank Channa Bambaradeniya (IUCN – The World Conservation Union) for reviewing the manuscript. We would also like to thank Mendis Wickramasinghe (SLHS–Sri Lanka Herpetological Society) and Kelum Manamendra-Arachchi (WHT) for sending valuable literature. Our heartfelt thanks go to the following persons for their support; Naalin Perera, Dilup

Chandranimal, Sarath Ekanayake, Sampath Goonatilake, Vimukthi Weeratunga, Prasanna Samarawickrama, Sandun Perera, Suranjan Fernando and Nalinda Peiris, Vimukthi Weerathunga, Pradeep Samarawickrama (IUCN – The World Conservation Union). Finally, we thank Chamila Soysa, Toshana Peiris, Panduka Silva, Asanka Udayakumara, Anushka Kumarasinghe, Dimuthu Wickramasinghe and Thasun Amarasinghe (YZA–Young Zoologists' Association) for their kind help during the field visit and other activities in Nilgala Forest Area.

LITERATURE CITED

- Bambaradeniya, C.N.B., M.S.J. Perera, W.P.N. Perera, L.J.M. Wickramasinghe, L.D.C.B. Kekulandala, V.A.P. Samarawickrema, R.H.S.S. Fernando, and V.A.M.P.K. Samarawickrema. 2003. Composition of faunal species in the Sinharaja world heritage site in Sri Lanka. *Sri Lanka Forester* 26:21-40.
- Bossuyt, F., M. Meegaskumbura, N. Beenaerts, D.J. Gower, R. Pethiyagoda, K. Roelants, A. Mannaert, M. Wilkinson, M.M. Bahir, K. Manamendra-rachchi, P.K.L. Ng, C.J. Schneider, O.V. Oommen, and M.C. Milinkovitch. 2004. Local endemism within the Western Ghats – Sri Lanka Biodiversity Hotspot. *Science* 306:479-481.
- Burgett, A.A., C.D. Wright, G.R. Smith, D.T. Fortune, and S.L. Johnson. 2007. Impact of ammonium nitrate on Wood Frog (*Rana sylvatica*) tadpoles: effects on survivorship and behavior. *Herpetological Conservation and Biology* 2:29-34.
- Bury, R.B. 2006. Natural history, field ecology, conservation biology, and wildlife management: Time to connect the dots. *Herpetological Conservation and Biology* 1:56-61.
- Camargo, J.L., and V. Kapos. 1995. Complex edge effects on soil moisture and microclimate in central Amazonian forest. *Journal of Tropical Ecology* 11:205-221.
- De Silva, A. 1996. The Herpetofauna of Sri Lanka: a brief review. Graphic Land, Kandy, Sri Lanka.
- De Silva, A. 2006. Current status of the Reptiles of Sri Lanka. Pp. 134-163 *In* Fauna of Sri Lanka: Status of Taxonomy, Research and Conservation. Bambaradeniya, C.N.B. (Ed.). The World Conservation Union, Colombo, Sri Lanka and Government of Sri Lanka.
- De Silva, A., A.M. Bauer, C.C. Austin, S. Goonewardena, Z. Hawke, and V. Vanneck. 2004. The diversity of Nilgala forest, Sri Lanka, with special reference to its herpetofauna. *Lyriocephalus* 5(1&2):164-182.
- Dutta, S.K., and K.N. Manamendra-Arachchi. 1996. The Amphibian Fauna of Sri Lanka. Wildlife Heritage Trust of Sri Lanka.

- Fernando, S.S., L.J.M. Wickramasingha, and R.K. Rodrigo. 2007. A new species of endemic frog belonging to the (?) genus *Nannophrys* Günther, 1869 (Anura: Dicroglossinae) from Sri Lanka. *Zootaxa* 1403:55–68.
- Giri, V., and N. Chaturvedi. 2001. Preliminary survey of the Herpetofauna in the Western Ghats region of Maharashtra. *Tigerpaper* 8:1-7.
- Glista, D.J., T.L. DeVault, and J.A. DeWoody. 2008. Vertebrate road mortality predominantly impacts amphibians. *Herpetological Conservation and Biology* 3:77-87.
- Gray, M.J., L.M. Smith, D.L. Miller, and C.R. Bursey. 2007. Influences of agricultural land use on *Clinostomum attenuatum* metacercariae prevalence in southern Great Plains amphibians, U.S.A. *Herpetological Conservation and Biology* 2:23-28.
- Gunatilleke, I.A.U.N., and C.V.S. Gunatilleke. 1990. Distribution of floristic richness and its conservation in Sri Lanka. *Conservation Biology* 4:21-31.
- Gunawardene, N.R., A.E.D. Daniels, I.A.U.N. Gunatilleke, C.V.S. Gunatilleke, P.V. Karunakaran, K.G. Nayak, S. Prasad, P. Puyravaud, B.R. Ramesh, K.A. Subramanian, and G. Vasanthi. 2007. A brief overview of the Western Ghats – Sri Lanka biodiversity Hotspot. *Current Science* 93:1567-1572.
- Hettige, U.S.B., L.J.M. Wickramasinghe, T.G.M. Priyadarshana, K. Gunawardena, L.I. Perera, and A. Manorathna. 2000. Fauna of Gal Oya National Park. *Sri Lanka Naturalist* 3:55-61.
- IUCN Sri Lanka. 2000. The 1999 Red List Threatened Fauna and Flora of Sri Lanka, IUCN Sri Lanka, Colombo.
- IUCN Sri Lanka and the Minister of Environmental Natural Resources (MENR), 2007. The 2007 Red List of Threatened Fauna and Flora of Sri Lanka. Colombo, Sri Lanka.
- Kapfer, J.M., M.B. Sandheinrich, and M.G. Knutson. 2007. Use of FETAX to examine acute survival of *Xenopus laevis* larvae in water from natural and constructed ponds in the Midwest. *Herpetological Conservation and Biology* 2:119-126.
- Karraker, N.E. 2007. Are embryonic and larval Green Frogs (*Rana clamitans*) insensitive to road deicing salt? *Herpetological Conservation and Biology* 2:35-42.
- Karunaratna, D.M.S.S. and A.A.T. Amarasinghe. 2007. Observation on the breeding behavior of *Philautus regius* Manamendra-Arachchi and Pethiyagoda, 2005 (Amphibia: Ranidae: Rhacophorinae) in Nilgala, Monaragala District in Sri Lanka. *Russian Journal of Herpetology* 14:133-136.
- Karunaratna, D.M.S.S., A.A.T. Amarasinghe, U.T.I. Abeywardena, M.D.C. Asela, and D.G.R. Sirimanna. 2006. Preliminary study on herpetofauna diversity of Nilgala forest area in Monaragala District, Sri Lanka. International Forestry and Environment Symposium, Department of Forestry and Environmental Science, University of Sri Jayewardenepura, Sri Lanka 11:74.
- Karunaratna, D.M.S.S., and D.M.G.N. Karunaratna. 2005. An unusual behavior of *Otocryptis nigristigma* Bahir and Silva, 2005 (Reptilia: Agamidae) observed at Nilgala forest in Sri Lanka. *Sri Lanka Naturalist* 7(1&2):21-22.
- Karunaratna, D.M.S.S., U.T.I. Abeywardena, M.D.C. Asela, D.M.G.N. Karunaratna, D.G.R. Sirimanna, and A.A.T. Amarasinghe. 2005. First record of the *Chamaeleo zeylanicus* Laurenti, 1768 (Reptilia: Chamaeleonidae) from the Nilgala forest in Sri Lanka. *Loris* 24(1&2):18-20.
- Knopf, A.A. 1998. Familiar Reptiles and Amphibians of North America, Chanticleer Press, New York, USA.
- Manamendra-Arachchi, K. 2000. Know your frogs. *Sri Lanka Nature* 2(5):4–16.
- Manamendra-Arachchi, K, and R. Pethiyagoda. 2005. The Sri Lankan shrub-frogs of the genus *Philautus* Gistel, 1848 (Ranidae: Rhacophorinae) with description of 27 new species. Pp. 163-303 *In* Contribution to Biodiversity Exploration and Research in Sri Lanka. Yeo, D.C.J., P.K.L. Ng, and R. Pethiyagoda. (Eds.). Raffles Bulletin of Zoology, Supplement No. 12.
- Manamendra-Arachchi, K, and R. Pethiyagoda. 2006. Sri Lankawe Ubhayajeeven “Amphibian Fauna of Sri Lanka” (text in Sinhala). Wildlife Heritage Trust of Sri Lanka.
- Manamendra-Arachchi, K, S. Batuwita, and R.Pethiyagoda. 2007. A taxonomic revision of the Sri Lanka day-geckos (Reptilia: Gekkonidae: *Cnemaspis*), with description of new species from Sri Lanka and Southern India. *Zeylanica* 7:9-122.
- McCallum, M.L. 2007. Amphibian decline or extinction? Current declines dwarf background extinction rate. *Journal of Herpetology* 41:483-491.
- McCallum, M.L., and J.L. McCallum. 2006. Publication trends of natural history and field studies in Herpetology. *Herpetological Conservation and Biology* 1:62-67.
- McCallum, M.L., and S.E. Trauth. 2003. A forty-three year museum study of Northern Cricket Frog (*Acris crepitans*) abnormalities in Arkansas: Upward trends and distributions. *Journal of Wildlife Diseases* 39:522-528.
- Meegaskumbura, M., F. Bossuyt, R. Pethiyagoda, K. Manamendra-Arachchi, M.M. Bahir, M.C. Milinkovitch, and C.J. Schneider. 2002. Sri Lanka: an amphibian hotspot. *Science* 298:379.
- Meegaskumbura, M., and K. Manamendra-Arachchi. 2005. Description of eight new species of shrub-frogs (Ranidae: Rhacophorinae: *Philautus*) from Sri Lanka. Pp. 305-338 *In* Contribution to Biodiversity Exploration and Research in Sri Lanka. Yeo, D.C.J.,

- P.K.L. Ng, and R. Pehiyagoda. (Eds.). Raffles Bulletin of Zoology, Supplement No. 12.
- Meegaskumbura, M., K. Manamendra-Arachchi, C.J. Schneider, and R. Pehiyagoda. 2007. New species amongst Sri Lanka's extinct shrub frogs (Amphibia: Rhacophoridae: *Philautus*). *Zootaxa* 1397:1-15.
- Mendelson III, J.R., K.R. Lips, R.W. Gagliardo, G.B. Rabb, J.P. Collins, J.E. Diffendorfer, P. Daszak, D.R. Ibanez, K.C. Zippel, D.P. Lawson, K.M. Wright, S.N. Stuart, C. Gascon, H.R. DA Silva, P.A. Burrowes, R.L. Joglar, E. La Marca, S. Lotters, L.H. Du Preez, C. Weldon, A. Hyatt, J.V. Rodriguezmahecha, S. Hunt, H. Robertson, B. Lock, C.J. Raxworthy, D.R. Frost, R.C. Lacy, R.A. Alford, J.A. Campbell, G. Parra-Olea, F. Bolanos, J.J.C. Domingo, T. Halliday, J.B. Murphy, M.H.Wake, L.A. Coloma, S.L. Kuzim, M.S. Price, K.M. Howell, M. Lau, R. Pehiyadoda, M. Boone, M.J. Lannoo, A.R. Blaustein, A. Dobson, R.A.L. Griffiths, M. Crump, D.B. Wake, and E.D. Brodie, JR. 2006. Confronting amphibian declines and extinctions. *Science* 313:48.
- Murcia, C. 1995. Edge effects in fragmented forests: Implications for conservation. *Trends in Ecology and Evolution* 10:58-62.
- Pehiyagoda, R., K. Manamendra-Arachchi, M.M. Bahir, and M. Meegaskumbura. 2006. Sri Lanka amphibians: Diversity, Uniqueness and Conservation. Pp. 125-133 *In* Fauna of Sri Lanka: Status of Taxonomy, Research and Conservation. Bambaradeniya, C.N.B. (Ed). The World Conservation Union, Colombo, Sri Lanka and Government of Sri Lanka.
- Pough, F.H., R.M. Andrews, J.E. Cadle, M.L. Crump, A.H. Savitzky, and K.D. Wells. 2004. *Herpetology*. 3rd Edition. Pearson Prentice Hall, San Francisco, USA.
- Samarawickrama, V.A.M.P.K., K.B. Ranawana, D.R.N.S. Rajapaksha, N.B. Ananjeva, N.L. Orlov, J.M.A.S. Ranasinghe, and V.A.P. Samarawickrama. 2006. A new species of the genus *Cophotis* (Squamata: Agamidae) from Sri Lanka. *Russian Journal of Herpetology* 13:207-214.
- Senanayake, F.R., M. Soule, and J.W. Senner. 1977. Habitat values and endemism in the vanishing rainforest of Sri Lanka. *Nature* 265:351-354.
- Tadevosyan, T.L. 2006. Habitat suitability for reptiles in the Goravan Sands Sanctuary, Armenia. *Herpetological Conservation and Biology* 1:39-44.
- Wickramasinghe L.J.M., and D.A.I. Munindradasa. 2007. Review of the genus *Cnemaspis* Strauch, 1887 (Sauria: Gekkonidae) in Sri Lanka, with the description of five new species. *Zootaxa* 1490:1-63.
- Wickramasinghe L.J.M., R.K. Rodrigo, N. Dayawansa, and U.L.D. Jayantha. 2007. Two new species of *Lancascincus* (Squamata: Scincidae) from Sripada Sanctuary (Peak Wilderness), in Sri Lanka. *Zootaxa* 1612:1-24.
- Wijesinghe, M.R., and P.N. Dayawansa. 2002. The amphibian fauna at two altitudes in the Sinharaja rainforest, Sri Lanka. *Herpetological Journal* 12:175-178.


APPENDIX. Species observed at Nilgala Forest Area, Monaragala District , Sri Lanka.



Pollonnaruwa Shrub Frog
(*Philautus regius*)



Ferguson's Shrub Frog
(*Philautus fergusonianus*)



Tribble Rock Frog
(*Nannophrys naeyakai*).



Common House Toad
(*Bufo melanostictus*).



Red Narrow Mouth Frog
(*Microhyla rubra*).



Chunam Tree Frog
(*Polypedates maculatus*).



Common Bull Frog
(*Kaloula taprobanica*).



Banded Sand Frog
(*Sphaerotheca breviceps*).



Sri Lanka Wood Frog
(*Rana gracilis*).



Common Hourglass Tree Frog
(*Polypedates cruciger*).



SURANJAN KARUNARATHNA has studied amphibian and reptile ecology, biology, behavior and conservation in Sri Lanka for over 10 years. During this time, he started numerous educational and awareness programs related to herpetofauna and their conservation. He is also an instructor in the reptile study group of Young Zoologists' Association of Sri Lanka, based at the National Zoological Gardens, and currently works as a project assistant for the IUCN, Sri Lanka Country office in Colombo. During his career he has worked with diverse groups including government agencies and non government organizations in Sri Lanka.



INDIKA ABEYWARDENA is an ecologist of protected area management project in the Department of Wildlife Conservation. He is an undergraduate and earned his diploma in Archaeology Heritage Management from the University of Kelaniya. He is also an instructor in the flora study group of the Young Zoologists' Association of Sri Lanka, based at the National Zoological Gardens, Sri Lanka. His hobbies include Nature and Wildlife Photography, Hiking and exploration, and Conservation of Sri Lankan Orchids. His interests span herpetology, invasive species, conservation of rare plants and plant ecology.



CHANDANA ASELA is an undergraduate in Science at The Open University of Sri Lanka, and earned a Diploma in biodiversity management from the University of Colombo. He is a senior Member of the Young Zoologists' Association of Sri Lanka, National Zoological Garden, Dehiwala. He has worked in the environmental field for 10 years and promotes environmental awareness, research and conservation. He has worked as an ecologist with the IUCN, Sri Lanka Country office, Colombo. His interests span avifauna, mammals and plant biogeography. He is currently a program officer in the United Nations Office for Project Services in Sri Lanka.



BHATHIYA KEKULANDALA is a researcher in biodiversity science with a keen interest in conservation and management of fragile biological resources of Sri Lanka. His research focuses on indigenous forest resource management systems and framework development for assessment of ecological services. He works in both an official and in a voluntary capacity with government and non-government sectors including the Department of Wildlife Conservation, Forestry Department, Department of National Zoological Gardens, Ministry of Environmental and Natural Resources, Customs of Sri Lanka (Biodiversity Unit), University of Colombo, IUCN, and the Young Zoologists' Association of Sri Lanka.