PREVALENCE OF SALMONELLA IN INTESTINAL MUCOSAL SAMPLES FROM FREE-RANGING RED-EARED SLIDERS (TRACHEMYS SCRIPTA ELEGANS) IN ILLINOIS

ANNE M. READEL^{1,2}, CHRISTOPHER A. PHILLIPS^{1,2}, AND TONY L. GOLDBERG³

¹Illinois Natural History Survey, Institute for Natural Resource Sustainability, University of Illinois, Champaign, Illinois 61820, USA

²Program in Ecology, Evolution, and Conservation Biology, University of Illinois, Urbana, Illinois 61801, USA

³Department of Pathobiological Sciences, School of Veterinary Medicine, University of Wisconsin, Madison,

Wisconsin 53706, USA, e-mail: readel@wisc.edu

Abstract.—Turtles can be carriers of bacteria of the genus *Salmonella*, and numerous studies have documented *Salmonella* shedding in both captive and free-living populations. Because turtles may shed *Salmonella* intermittently, however, data based on fecal samples may underestimate the true prevalence of the bacterium in turtles. We examined intestinal mucosal scrapings using molecular methods to identify *Salmonella* in free-ranging Red-eared Sliders (*Trachemys scripta elegans*) to determine carriage rates. *Salmonella* was detected in 11% (8/73) of turtles. Prevalence of infection did not vary among the nine ecologically varied ponds sampled. The prevalence of *Salmonella* infection in this study was higher than documented in a previous study that was conducted on conspecifics in the same area and during the same year but using fecal samples, suggesting that free-ranging Red-eared Sliders can harbor *Salmonella* but not shed the bacterium in feces.

Key Words.-bacteria; infections; Red-eared Sliders; Salmonella; Trachemys scripta; turtles

INTRODUCTION

Reptiles are generally asymptomatic carriers of bacteria in the genus Salmonella (Chiodini and Sundberg 1981) and have been identified as a source of infection in human cases and outbreaks of salmonellosis (Chiodini and Sundberg 1981; Mermin et al. 2004; Harris et al. 2009). The association between salmonellosis and pet turtles motivated a 1975 federal ban on the sale of turtles with a carapace of < 10.2 cm (4 in) in the United States (Cohen et al. 1980). This ban prevented an estimated 100,000 cases of turtle-associated salmonellosis in children between 1970 and 1976 alone (Cohen et al. 1980). These public health concerns have also prompted studies to identify the prevalence of Salmonella in captive turtles (Otis and Behler 1973; Siebeling et al. 1984; Shane et al. 1990; Abalem de Sá and Solari 2001). More recently, studies of free-ranging turtles have reported prevalence of shed Salmonella as low as 0% (Richards et al. 2004; Saelinger et al. 2006; Readel et al. 2008) and as high as 50-100% (Chambers and Hulse 2006; Hahn et al. 2007; Gaertner et al. 2008a), highlighting the extreme variability in shedding rates characteristic of these bacteria.

In a recent study, we failed to isolate *Salmonella* from cloacal samples collected from Red-eared Sliders (*Trachemys scripta elegans*) living in multiple southern Illinois ponds that varied in pond type (natural vs.

manmade), size, and biotic communities (Readel et al. 2008). Studies of cloacal shedding rates may provide an incomplete picture of the prevalence of *Salmonella* in free-ranging turtles because *Salmonella* shedding might be intermittent and stress-related (DuPonte et al. 1978; Burnham et al. 1998). Therefore, the prevalence of cloacal shedding could underestimate the true infection rate of turtles. Nevertheless, few studies have attempted to quantify carriage rates of *Salmonella* in free-ranging turtles (for an exception, see Saelinger et al. 2006), which limits not only accurate assessments of public health risk, but also our ability to examine the influence of host and environmental characteristics on rates of infection.

The aim of this study was to determine the prevalence Salmonella carriage rates by examining of gastrointestinal mucosal samples from free-ranging Redeared Sliders in southern Illinois. Red-eared Sliders are common in the pet trade, notorious as a globally invasive species, and inhabit a variety of aquatic habitats. Numerous studies have already quantified Salmonella shedding rates in this species in captivity (McCoy and Seidler 1973; Otis and Behler 1973; Chassis et al. 1986; Abalem de Sá and Solari 2001; Pasmans et al. 2002a); our study provides complementary information on rates of carriage independent of shedding. Additionally, because environmental and host life-history characteristics may influence Salmonella shedding in



FIGURE 1. The location of the study ponds where we captured Red-eared Sliders (Trachemys scripta elegans) in southeastern Illinois.

free-ranging turtles (Hidalgo-Vila et al. 2006; Gaertner et al. 2008a, b; Readel et al. 2008), we also examined the influence of host characteristics (body size, and body condition) and pond location on *Salmonella* infection rates.

MATERIALS AND METHODS

We captured turtles using baited hoop traps from nine water bodies in southern Illinois from 2-25 August 2006 (Fig. 1; Table 1), a region characterized primarily by a mosaic landscape of forest, grassland, and intensive row crop agriculture. Ponds varied in size (2–50 hectares), and in the percentage of intensive row crop agriculture (2–50%) and forest (0–54%) surrounding each pond edge within a 125 m buffer. Half of the ponds were natural oxbow lakes along the Ohio River floodplain that supported a high diversity of fishes as well as diverse turtle assemblages (Dreslik et al. 2005). The others were manmade ponds created in or before the 1970's as city reservoirs or from highway borrow pits, contained only common lentic turtle species, and some have been

stocked with game fish. All ponds were used for recreational fishing. Turtle abundance, defined as the number of individual turtles of any species captured per trap hour, was also determined for each pond.

We collected only adult male turtles for this study. This was done to minimize potential life history differences among individuals and because this study was conducted as part of a larger investigation that required only males. We handled turtles using individual bleach-decontaminated plastic tubs and instruments, and we used sterile procedures throughout the study to prevent cross-contamination. Upon return to the laboratory, we measured turtles for plastron length (PL) to the nearest mm and we killed them by overexposure to halothane vapors for two hours. Halothane induces anesthesia rapidly and is the most effective and preferred inhalant anesthetic for euthanasia of small animals (American Veterinary Medical Association. 2007. AVMA Guidelines on Euthanasia. Available from http://www.avma.org/resources/ euthanasia.pdf [Accessed 2 July 2010]). Death was confirmed by prolonged (in excess of 30 min) lack of

TABLE 1. Prevalence of *Salmonella* in male Red-eared Sliders (*Trachemys scripta elegans*) from nine ecologically varied ponds in Illinois. Confidence intervals (95%) for the prevalence of *Salmonella* in turtles were calculated using the modified Wald method (Agresti and Coull, 1998).

Pond	Location	No. Turtles Sampled	Prevalence of Salmonella (%)	95% Confidence Interval
300N	37.9229N, 88.0992W	8	12.5	0.1 - 49.2
Big	37.7258N, 88.1162W	8	12.5	0.1 - 49.3
Calvin	37.7371N, 88.5118W	7	28.6	7.5 - 64.8
Fish	37.7228N, 88.1043W	8	12.5	0.1 - 49.3
Flat	38.1136N, 88.0022W	10	0.0	0.0 - 32.1
Hulda	37.7841N, 88.1034W	7	14.3	0.5 - 53.4
Norris	37.9810N, 88.3139W	10	0.0	0.0 - 32.1
Omaha	37.8987N, 88.3253W	8	12.5	0.1 - 49.3
Smugglers	37.7367N, 88.5215W	7	14.3	0.5 - 53.4
Total		73	11.0	5.4 - 20.4

righting response. Turtles were then transferred to a -80°C freezer until dissection in 2008. Turtles were dissected, and a mucosal scrape was obtained from the small and large intestine of each turtle. Scrapes were collected by scraping the length of the mucosal lining with the blunt end of a pair of sterile forceps after large food particles and parasites had been removed. The amount of mucus collected from each scrape varied between < 50 µl and 2 mL. Material from scrapes was placed in 2 mL microcentrifuge tubes and stored at -80°C. Additionally, all body fat was removed from each turtle and weighed to the nearest 0.01 g on an electronic balance (Denver Instrument Company, Model XE-100A, Denver, Colorado, USA) for a direct estimate of body condition (fat weight / turtle PL).

We subjected mucosal samples to a diagnostic PCR to identify Salmonella. We extracted DNA from 50-100 µl of intestinal mucus using the ZR-96 Genomic DNA Kit (Zymo Research, Orange, California, USA) eluted in a 50 µl volume, according to the manufacturer's protocol. The Salmonella specific primer pair invA-1 (5'-ACAGTGCTCGTTTACGACC TGAAT) and invA-2 (5'- AGACGACTGGTACTGAT CGATAAT) was used to amplify a 243 bp segment of the Salmonella invA gene, which has been shown to be both sensitive and specific for this bacterial genus (Chiu and Ou 1996). As an internal control for DNA extraction and amplification, we used another published PCR primer pair of 16S-F (5'-AGACTGCTACGGGAG GCAGCAGT) and 16S-R (5'- GTTGCGCTCGTTGC GGGACTTAA) to amplify a 755 bp fragment of the 16S ribosomal RNA subunit gene, which is present in all bacteria (Villalobo and Torres 1998). PCR was conducted in a 25 µl volume using the FailSafe[™] PCR System (Epicentre Biotechnologies, Madison. Wisconsin, USA) containing 12.5 µl FailSafe™ PCR PreMix Buffer B, 0.6 µM each of primer, 0.5 µl FailSafeTM Enzyme Mix, and 1 µl Template DNA. Reactions were cycled in an i-Cycler thermocycler (BioRad Inc, Hercules, California, USA) at 94°C for 2 min, then for 35 cycles at 94°C for 30 s, 57°C for 30 s,

and 72°C for 45 s, followed by a final 72°C 4 min extension step and an indefinite 4°C soak. We used a sample of wild-type *Salmonella enterica* serotype Typhimurium DNA from the University of Illinois Veterinary Diagnostic Laboratory as a positive control, and we used sterile distilled water as a negative control. Amplicons were visualized by electrophoresis in 1.5% agarose /Tris-acetate-EDTA (TAE) buffer followed by ethidium bromide staining and digital imaging under ultraviolet light. We considered samples positive only if PCR produced both the 243 bp *invA* band and the 755 bp internal control 16S rRNA band.

To determine the sensitivity of our PCR, we used a NanoDropTM spectrophotometer (Thermo Scientific, Waltham, Massachusetts, USA) to identify the concentration (ng/µl) of DNA in our positive control. The number of genome equivalents per sample was then determined from the concentration of DNA in this sample and the known size of the *Salmonella* genome. We then subjected serial dilutions from $10^5 - 10^0$ genome equivalents per reaction to our PCR protocol, and we determined the sensitivity of the PCR as the lowest concentration at which both the 243 bp *invA* band and the 755 bp internal control 16S rRNA band could be detected.

We fitted a generalized linear model to a binomial distribution to examine the effect of host size and body condition on prevalence of *Salmonella* in turtles. To determine the differences in *Salmonella* prevalence among ponds, we used a Fisher's Exact Test. Confidence intervals (95%) for the prevalence of *Salmonella* spp. in turtles were calculated using the modified Wald method (Agresti and Coull 1998). We used $\alpha = 0.05$ to test for significance in all cases. Statistics were performed using the computer program SAS (SAS Institute, Cary, North Carolina, USA).

RESULTS

We captured 73 free-ranging male Red-eared Sliders that were between 123–210 mm PL with body

conditions between 0.00-0.20 g fat/mm PL from 2-25 August 2006 (Table 1). We dissected all turtles and then collected 146 mucosal scrapes from the 73 turtles. However, the small or large intestine sample was $< 50 \,\mu$ l for five turtles. For those turtles, we pooled the small and large intestine mucosal scrapes to obtain a sufficient quantity of mucus for molecular analyses. Therefore, we analyzed 141 mucosal samples and eight individuals tested positive for Salmonella (two individuals tested positive for both mucosal scrapes, six individuals tested positive from the large intestine sample only). The prevalence of Salmonella spp. in the male turtles sampled was therefore 11% (8/73) with a 95% confidence limit of 5.4-20.4% for the entire population of 73 turtles (Table 1). Among the nine ponds sampled, the prevalence of Salmonella spp. in male turtles ranged between 0% and 28.6% (Table 1). Our PCR protocol resulted in a sensitivity of detection of 10^2 to 10^3 genome equivalents per reaction. Because we did not isolate Salmonella, we were unable to determine bacterial serotypes. The prevalence of Salmonella spp. in turtles did not vary significantly with turtle size ($\chi^2 = 2.13$, df = 1, P = 0.145), turtle body condition ($\chi^2 = 0.81$, df = 1, P = 0.368), or pond (P = 0.699, Fisher's Exact Test).

DISCUSSION

The prevalence of Salmonella in the male turtles in our study was 11% overall (CI: 5.4-20.4%) and ranged from 0% to 28.6% among the nine ponds sampled. To our knowledge, our study is the first to examine gastrointestinal mucosal samples from free-ranging turtles to determine Salmonella carrying rates. One other study examined 16 gastrointestinal mucosal samples from free-ranging turtles submitted to a clinic, but did not find Salmonella (Saelinger et al. 2006). Information on Salmonella carriage rates is important because Salmonella shedding in reptiles may be intermittent and stress-induced (DuPonte et al. 1978; Burnham et al. 1998), such that studies of Salmonella shedding could underestimate the true prevalence of infection. The results of our study indirectly support this possibility because the prevalence of Salmonella in male Red-eared Sliders was higher than that reported in a similar study of Red-eared Slider shedding rates (Readel et al. 2008). This study was conducted in the same region and during the same year as our study, which did not detect Salmonella in any of the 100 T. scripta sampled using cloacal swabs, which were analyzed using a combination of culture and molecular techniques (95% confidence interval of Salmonella prevalence was 0-4.4%, Readel et al. 2008). The fact that different methods were used between the studies makes comparisons of prevalence difficult, however, because detection methods can vary in their sensitivity (Mitchell et al. 2000). If the methods used to identify Salmonella

from cloacal swabs were less sensitive than those used on gastrointestinal mucosal samples, they may have failed to detect low-level infections and intermittent shedding, and may have accounted for the differences in prevalence between the two studies.

Variation in the sensitivity of different detection methods may also partially explain the wide range of Salmonella prevalence reported in other studies (Gaertner et al. 2008a; see below). Although our results were consistent with many other studies that have reported a moderate-low prevalence of Salmonella in free-ranging chelonians (8%, Jackson et al. 1969; 5%, Harwood et al. 1999; 0%, Richards et al. 2004; 0%, Saelinger et al. 2006; 5%, Lockhart et al. 2008; 0%, Readel et al. 2008; 10%, Charles-Smith et al. 2009), they were considerably lower than those of many other studies (32%, Briones et al. 2004; 100%, Chambers and Hulse 2006; 50%, Hahn et al. 2007; 51%, Gaertner et al. 2008a; 46%, Gaertner et al. 2008b), including other data on Red-eared Sliders (48%, Gaertner et al. 2008b). The relatively low prevalence of Salmonella in the male turtles in this study could be due to the sensitivity of our methods. Because the density of Salmonella organisms may be low in free-ranging turtles, highly sensitive procedures may be required to identify infections (Gaertner et al. 2008a). In other studies, a preenrichment and/or enrichment step were used to detect Salmonella from cloacal, carapace, and environmental samples (Hahn et al. 2007; Gaertner et al. 2008a; Charles-Smith et al. 2009), and may have been useful for identifying turtles with low levels of infection in this study. Nevertheless, we directly sampled portions of the gastrointestinal tract where Salmonella is known to reside (Pasmans et al. 2002b), we performed duplicate reactions that would have increased our probability of detection, we optimized our PCR for sensitivity, and we were able to isolate Salmonella.

Environmental characteristics may also increase Salmonella exposure or shedding rates in free-ranging turtles. Salmonella survives for long periods of time in the aquatic environment (Morse and Duncan 1974; Chiodini and Sundberg 1981) and ingestion of fecal material or contaminated water is a likely mode of infection (Chiodini and Sundberg 1981; Polo et al. 1999; Baudart et al. 2000). Therefore, a high number of turtles in closed systems, and consequent reinoculation of Salmonella through feces, is believed to be partially responsible for high levels of Salmonella in captive turtles (Kaufman et al. 1972; Chiodini and Sundberg 1981). The lower prevalence of *Salmonella* in the male turtles in our study suggests that environmental contamination of our ponds with Salmonella may be low, and possibly that turtle abundance in our ponds might not have represented overcrowding.

Additionally, we found no differences in *Salmonella* infection prevalence among ponds, despite a range of

pond characteristics that could have enhanced Salmonella infection or transmission. For instance, intensive row crop agriculture adjacent to ponds might have been expected to result in higher nutrient content in the pond water, promoting the growth of coliform bacteria, or forests adjacent to ponds might have had higher densities of animal reservoirs of Salmonella. The lack of an association between Salmonella prevalence and environmental characteristics is, however, consistent with the results of other studies. In those studies, the prevalence of Salmonella in free-ranging turtles was similar across seven sites along one river (range: 40-75%, Gaertner et al. 2008a), and between spring (53%, Hahn et al. 2007) and slough (50%, Gaertner et al. 2008b) headwater sections of a river. These results suggest that Salmonella infection rates in free-ranging aquatic turtles are stable across microhabitats within ecosystems.

Although our study did not find associations among host and environmental characteristics and Salmonella infection levels, our study was cross-sectional and did not examine temporal variability in infection rates. Furthermore, we only sampled male turtles; however, this is unlikely to have had a large impact on the Salmonella prevalence reported in this study because other studies have not found Salmonella prevalence to differ between male and female turtles (Gaertner et al. 2008a: Lockhart et al. 2008). Turtles were also only sampled from southeastern Illinois, so the generality of our findings to wild turtle populations across the state and elsewhere may be limited. Additional longitudinal sampling, however, might shed light on the underlying factors that explain Salmonella infection rates in freeranging turtles. Finally, our results emphasize that people working with or interacting with wild turtles need to be aware of the possibility of Salmonella infection and should use precautions when handling turtles.

Acknowledgments.—This research was supported by the Illinois Department of Natural Resources Wildlife Preservation Fund, Sigma Xi, the Chicago Herpetological Society, the Illinois State Academy of Sciences, the R. Weldon Larimore/Jordan Creek Endowment Fund, the Minnesota Herpetological Society, and the University of Illinois Program in Ecology, Evolution, and Conservation Biology. The authors thank Festus Lee for logistic assistance in the field, Traci Tranby-McLachlan for assistance with optimizing our PCR protocol, Mark Mitchell for demonstrating turtle necropsy techniques, and Amanda Wolff, Ursula Sieklucki, Adrien Nickel, Elsa Holden, Heather Davis, Kim Wangen, and Natalia Piejko for general assistance in the laboratory. All research and euthanasia procedures were conducted in accordance University of Illinois, Urbana-Champaign with Institutional Animal Care and Use Committee protocol

#06171, and turtles were deposited in the Illinois Natural History Survey Amphibian and Reptile Collection after the study was completed.

LITERATURE CITED

- Abalem de Sá, I.V., and C.A. Solari. 2001. *Salmonella* in Brazilian and imported pet reptiles. Brazilian Journal of Microbiology 32:293–297.
- Agresti, A., and B.A. Coull. 1998. Approximate is better than "exact" for interval estimation of binomial proportions. American Statistician 52:119–126.
- Baudart, J., K. Lemarchand, A. Brisabois, and P. Lebaron. 2000. Diversity of *Salmonella* strains isolated from the aquatic environment as determined by serotyping and amplification of the ribosomal DNA spacer regions. Applied and Environmental Microbiology 66:1544–1552.
- Briones, V., S. Téllez, J. Goyache, C. Ballesteros, M. del Pilar Lanzarot, L. Dominguez, and J.F. Fernández-Garayzábal. 2004. *Salmonella* diversity associated with wild reptiles and amphibians in Spain. Environmental Microbiology 6:868–871.
- Burnham, B.R., D.H. Atchley, R.P. DeFusco, K.E. Ferris, J.C. Zicarelli, J.H. Lee, and F.J. Angulo. 1998. Prevalence of fecal shedding of *Salmonella* organisms among captive Green Iguanas and potential public health implications. Journal of the American Veterinary Medical Association 213:48–50.
- Chambers, D.L., and A.C. Hulse. 2006. *Salmonella* serovars in herpetofauna of Indiana County, Pennsylvania. Applied and Environmental Microbiology 72:3771–3773.
- Charles-Smith, L.E., G.A. Lewbart, M.J. Aresco, and P. Cowen. 2009. Detection of *Salmonella* in Gopher Tortoises (*Gopherus polyphemus*) during two relocation efforts in Florida. Chelonian Conservation and Biology 8:213–216.
- Chassis G., E.M. Gross, Z. Greenberg, M. Tokar, N. Platzner, R. Mizrachi, and A. Wolff. 1986. *Salmonella* in turtles imported to Israel from Louisiana. Journal of the American Medical Association 256:1003.
- Chiodini, R.J., and J.P. Sundberg. 1981. Salmonellosis in reptiles: a review. American Journal of Epidemiology 113:494–499.
- Chiu, C.H., and J.T. Ou. 1996. Rapid identification of *Salmonella* serovars in feces by specific detection of virulence genes, *invA* and *spvC*, by an enrichment broth culture-multiplex PCR combination assay. Journal of Clinical Microbiology 34:2619–2622.
- Cohen M.L., M. Potter, R. Pollard, and R.A. Feldman. 1980. Turtle-associated salmonellosis in the United States: effect of public health action, 1970 to 1976. Journal of the American Medical Association 243:1247–1249.

- Dreslik, M.J., A.R. Kuhns, and C.A. Phillips. 2005. Structure and composition of a southern Illinois freshwater turtle assemblage. Northeastern Naturalist 12:173–186.
- DuPonte, M.W., R.M. Nakamura, and E.M.L. Chang. 1978. Activation of latent *Salmonella* and *Arizona* organisms by dehydration in Red-eared Turtles, *Pseudemys scripta elegans*. American Journal of Veterinary Research 39:529–530.
- Gaertner, J.P., D. Hahn, J. Jackson, M.R.J. Forstner, and F.L. Rose. 2008a. Detection of Salmonellae in captive and free-ranging turtles using enrichment culture and polymerase chain reaction. Journal of Herpetology 42:223–231.
- Gaertner, J.P., D. Hahn, F.L. Rose, and M.R.J. Forstner. 2008b. Detection of Salmonellae in different turtle species within a headwater spring ecosystem. Journal of Wildlife Diseases 44:519–526.
- Hahn, D., J.P. Gaertner, M.R.J. Forstner, and F.L. Rose. 2007. High resolution analysis of salmonellae from turtles within a headwater spring ecosystem. FEMS Microbiology Ecology 60:148–155.
- Harris, J.R., D. Bergmire-Sweat, J.H. Schlegel, K.A. Winpisinger, R.F. Klos, C. Perry, R.V. Tauxe, and M.J. Sotir. 2009. Multistate outbreak of *Salmonella* infections associated with small turtle exposure, 2007– 2008. Pediatrics 124:1388–1394.
- Harwood, V.J., J. Butler, D. Parrish, and V. Wagner. 1999. Isolation of fecal coliform bacteria from the Diamondback Terrapin (*Malaclemys terrapin centrata*). Applied and Environmental Microbiology 65:865–867.
- Hidalgo-Vila, J., C. Díaz-Paniagua, C. de Frutos-Escobar, C. Jiménez-Martínez, and N. Pérez-Santigosa. 2006. *Salmonella* in free living terrestrial and aquatic turtles. Veterinary Microbiology 119:311–315.
- Jackson, M.M., C.G. Jackson, and M. Fulton. 1969. Investigation of the enteric bacteria of the Testudinata –I: Occurrence of the genera *Arizona, Citrobacter, Edwardsiella,* and *Salmonella.* Journal of Wildlife Diseases 5:328–329.
- Kaufman, A.F., M.D. Fox, G.K. Morris, B.T. Wood, J.C. Feeley, and M.K. Frix. 1972. Turtle-associated salmonellosis. III. The effects of environmental salmonellae in commercial turtle breeding ponds. American Journal of Epidemiology 95:521–528.
- Lockhart, J.M., G. Lee, J. Turco, and L. Chamberlin. 2008. Salmonella from Gopher Tortoises (Gopherus polyphemus) in South Georgia. Journal of Wildlife Diseases 44:988–991.
- McCoy, R.H., and R.J. Seidler. 1973. Potential pathogens in the environment: isolation, enumeration, and identification of seven genera of intestinal bacteria

associated with small green pet turtles. Applied Microbiology 25:534–538.

- Mermin, J., L. Hutwagner, D. Vugian, S. Shallow, P. Daily, J. Bender, J. Koehler, R. Marcus, and F.J. Angulo. 2004. Reptiles, amphibians, and human *Salmonella* infection: a population-based, case-control study. Clinical Infectious Diseases 38:253–261.
- Mitchell, M.A., S.M. Shane, K. Orr, J. Nevarez, K. Maurer, D. Pesti, S. Sanchez, R.E. Wooley, and B. Ritchie. 2000. *Salmonella* diagnostic testing in the absence of a gold standard. Proceedings of the Association of Reptilian and Amphibian Veterinarians 7:143–144.
- Morse, E.V., and M.A. Duncan. 1974. Salmonellosis an environmental health problem. Journal of the American Veterinary Medical Association 165:1015–1019.
- Otis, V.S., and J.L. Behler. 1973. The occurrence of Salmonellae and *Edwardsiella* in turtles of the New York Zoological Park. Journal of Wildlife Diseases 9:4–6.
- Pasmans, F., P. De Herdt, and F. Haesebrouck. 2002a. Presence of *Salmonella* infections in freshwater turtles. Veterinary Record 150:692–693.
- Pasmans, F., P. De Herdt, J. Dewulf, and F. Haesebrouck. 2002b. Pathogenesis of infections with *Salmonella enterica* subsp. *Enterica* serovar Muenchen in the turtle *Trachemys scripta scripta*. Veterinary Microbiology 87:315–325.
- Polo, F.M., J. Figueras, I. Inza, J. Sala, J.M. Fleisher, J. Guarro. 1999. Prevalence of *Salmonella* serotypes in environmental waters and their relationships with indicator organisms. Antonie van Leeuwenhoek 75:285–292.
- Readel, A.M., C.A. Phillips, and T.L. Goldberg. 2008. Absence of cloacal shedding of *Salmonella* in wild Red-eared Sliders (*Trachemys scripta elegans*). Herpetological Review 39:427–430.
- Richards, J.M., J.D. Brown, T.R. Kelly, A.L. Fountain, and J.M. Sleeman. 2004. Absence of detectable *Salmonella* cloacal shedding in free-living reptiles on admission to the wildlife center of Virginia. Journal of Zoo and Wildlife Medicine 35:562–563.
- Saelinger, C.A., G.A. Lewbart, L.S. Christian, and C.L. Lemons. 2006. Prevalence of *Salmonella* spp in cloacal, fecal, and gastrointestinal mucosal samples from wild North American turtles. Journal of the American Veterinary Medical Association 229:266–268.
- Shane, S.M., R. Gilbert, and K.S. Harrington. 1990. *Salmonella* colonization in commercial pet turtles (*Pseudemys scripta elegans*). Epidemiology and Infection 105:307–316.

Herpetological Conservation and Biology

- Eradication of Salmonella and Arizona species from turtle hatchlings produced from eggs treated on commercial turtle farms. Applied and Environmental Microbiology 47:658-662.
- Siebeling, R.J., D. Caruso, and S. Neuman. 1984. Villalobo, E., and A. Torres. 1998. PCR for detection of Shigella spp. in mayonnaise. Applied and Environmental Microbiology 64:1242-1245.



ANNE READEL is a law student at the University of Wisconsin School of Law in Madison, WI. She received her Ph.D. from the University of Illinois in 2009 where she also received her B.S. in Animal Sciences in 2003. Her dissertation research focused on the effects of habitat on aquatic turtle health. In 2007, she worked at the National Oceanic and Atmospheric Administration (NOAA) in Washington, DC as a John Knauss Marine Policy Fellow. During law school, she has held legal clerkships at the U.S. Department of Justice, Environment and Natural Resources Division, and the U.S. Forest Service Patent Office. She also serves on the University of Wisconsin Lakeshore Nature Preserve Committee. She is interested in law and environmental policy. (Photographed by Tony L. Goldberg)



CHRISTOPHER A. PHILLIPS is a Research Program Leader at the Illinois Natural History Survey, Champaign. He received his Ph.D. from Washington University St. Louis in 1989 and his B.Sc. from Eastern Illinois University in 1983. His current research interests are in the fields of ecology and population genetics. Current questions focus on North American amphibians and reptiles. He is especially interested in population structure of wide ranging species and population viability. (Photographed by Kevin Cummings)



TONY GOLDBERG is a Professor of Epidemiology in the Department of Pathobiological Sciences at the University of Wisconsin-Madison. He received his D.V.M. from University of Illinois in 2000, his Ph.D. from Harvard University in 1996, and his B.A. from Amherst College in 1990. His research focuses on the ecology, epidemiology and evolution of infectious disease, combining field and laboratory studies to understand how pathogens in dynamic ecosystems around the world are transmitted among hosts, across complex landscapes, and over time. Dr. Goldberg holds additional appointments at the University of Wisconsin in the Nelson Institute for Environmental Studies, the Zoology Department, the Department of Population Health Sciences, and the Limnology and Marine Science Program, and is an Honourary Lecturer in Zoology at Makerere University, Uganda. (Photographed by Anne M. Readel)