GAMBUSIA HOLBROOKI PREDATION ON PSEUDACRIS FERIARUM TADPOLES

MARK STANBACK

Department of Biology, Davidson College, Davidson, North Carolina 28035, USA, e-mail: mastanback@davidson.edu

Abstract.—Although introduced world-wide for mosquito control, the Eastern Mosquitofish (*Gambusia holbrooki*) is now a major predator of amphibian larvae where it has been introduced. Because few researchers have examined the effects of *Gambusia* predation on anurans with which it has co-evolved, I compared survival of Upland Chorus Frog (*Pseudacris feriarum*) tadpoles in the presence and absence of *G. holbrooki*. I added 40 hatchling tadpoles to 12 cattle tanks, six of which contained *Gambusia*. After two weeks I found no tadpoles in tanks containing *Gambusia*, but many tadpoles in all the fish-free tanks. These results indicate that *Gambusia* are highly effective predators on small anuran larvae, even on species that share a long co-evolutionary history.

Key Words.—Eastern Mosquitofish; Gambusia holbrooki; predation; Pseudacris feriarum; tadpole; Upland Chorus Frog

INTRODUCTION

Introduced predators are one source of local amphibian declines and even extinctions (Stebbins and Cohen 1995; Hecnar and M'Closkey 1997; Kats and Ferrer 2003; Kiesecker 2003). In California, introduced larvae-eating fish have driven the Mountain Yellowlegged Frog (Rana muscosa) and the Red-legged Frog (R. aurora draytonii) from their optimal habitat (Bradford et al. 1993; Fisher and Shaffer 1996; Knapp and Matthews 2000). Introduced fish also negatively impacted the European Tree Frog (Hyla arborea) in Sweden (Brondmark and Edenhamn 1994). Members of the genus Gambusia have been especially problematic around the globe. Humans have introduced these "mosquitofish" for their purported ability to control numbers of mosquito larvae. However, researchers now know that introduced Gambusia eat more than just mosquito larvae (Miura et al. 1979; Bence 1988); amphibian larvae are clearly part of their diet (Webb and Joss 1997; Goodsell and Kats 1999; Lawler et al. 1999; Komak and Crossland 2000).

Researchers often assume that a lack of evolutionary contact with such predators is a primary reason these impacts are so devastating (Kats and Ferrer 2003). Many amphibians coexist naturally with the Eastern Mosquitofish (*Gambusia holbrooki*) in the southeastern United States. Consequently, researchers believe that such species have had the opportunity to evolve defensive mechanisms that allow for such coexistence (Kruse and Francis 1977; Petranka 1983). Yet few researchers have investigated the effects of *G. holbrooki* on amphibians in its native range (but see Grubb 1972; Baber and Babbit 2004). I used an experimental approach to assess the effects of the presence of *G. holbrooki* on the survivorship of young larvae of the Upland Chorus Frog (*Pseudacris feriarum*), a species

that breeds in both ephemeral and more permanent waters (Moriarty and Lannoo 2005).

MATERIALS AND METHODS

In late August 2005, I filled each of 12 oval 100gallon (378-1) cattle tanks with 375 1 of water, approximately 0.5 kg pine needles, and 250 mg alfalfa pellets. I arranged the tanks in two rows (six pairs of tanks) inside a chain-link fence near the Davidson College greenhouse in Davidson, Mecklenburg County, North Carolina (lat 35.4888, long -80.8275). One week later I inoculated each tank with 1 l of pond water. In late November, I captured wild G. holbrooki from a wetland area on the Catawba College ecological preserve in Salisbury, Rowan County, North Carolina (lat 35.6921, long -80.4865), a location also known as a breeding site for P. feriarum. The same day, I added 20 Gambusia (all between 2 and 4 cm in length) to one tank in each of the six pairs of tanks. I created similar size distributions of Gambusia in each of the six tanks. On 15 February 2006, I added an upright 9-cm x 19-cm x 39.5-cm cement cinderblock to each tank. On each cinderblock I placed a white 28 cm diameter ceramic dinner plate approximately 10 cm under the surface of the water.

On 19 February 2006 I collected ready-to-hatch egg masses of *P. feriarum* from a site in Rowan County (lat 35.6711, long -80.5729) and brought them into the laboratory in Davidson. By 20 February 2006, most of the eggs had hatched. At 1300 on 20 February, I placed 40 hatchling tadpoles into 500 ml plastic containers floating in each of the 12 cattle tanks. On 20 February 2000, I released 40 temperature-acclimated tadpoles into each of the tanks. Because *Gambusia* are primarily diurnal (Barrier and Hicks 2006), I introduced the tadpoles to the tanks at night to minimize initial predation. All tanks possessed refugia (pine needles,

algal growth) in which tadpoles could hide from fish. I conducted no water quality assessments. To assess survival of the tadpoles, I sampled each tank by counting the number of tadpoles visible on the white plate two weeks later, at 2000 on 5 March 2006. I compared the distribution of tadpoles in each pair of tanks using a Wilcoxon Signed-Rank test (two-tailed, $\alpha = 0.05$).

RESULTS

The presence of *Gambusia* had a clear effect on the survivorship of *P. feriarum* tadpoles (W = 10.5, df = 5, P = 0.031). In all the tanks without *Gambusia*, I observed at least two tadpoles on the submerged plate; I observed no tadpoles on the plates in the tanks containing *Gambusia* (Table 1). Moreover, close inspection of the walls and bottoms of the tanks revealed numerous additional tadpoles in the tanks without *Gambusia*; I detected no tadpoles in any of the tanks containing *Gambusia*. In subsequent weeks, I never observed tadpoles in fish-free tanks.

DISCUSSION

The most parsimonious explanation for the disappearance of tadpoles from tanks containing Gambusia is that the fish devoured most, if not all, of the tadpoles very quickly. Although I cannot rule out the possibility that the tadpoles engaged in cannibalism (see McDiarmid and Altig 1999; McCallum and Trauth 2001), Gambusia predation presumably explains the dramatic difference between fish-free and fish-filled tanks. This finding corresponds with recent research regarding the predatory effects of Gambusia. Many reports of predation of Australian tadpoles by introduced Gambusia exist (Morgan and Buttermer 1996; Komak and Crossland 2000; Hamer et al. 2002). Moreover, there is now experimental evidence detailing the precise nature of these negative impacts: Gamradt and Kats (1996) documented decreased survivorship of larval California Newts (Taricha torosa) in the presence of

Western Mosquitofish (*G. affinis*). Goodsell and Kats (1999) found similar decreases in survivorship of larval Pacific Tree Frogs (*Hyla regilla*). Lawler et al. (1999) found that *G. affinis* had multiple negative effects on California Red-legged Frog (*Rana aurora*) tadpoles: increased rates of tail injury, reduced metamorph size, decreased activity levels, and reduced rates of metamorphosis.

Multiple factors may influence the severity of Gambusia predation on anuran tadpoles, including the nutritional status of the Gambusia (Webb and Joss 1997; Pyke and White 2000), the movement of the tadpoles (Pyke and White 2000), and tadpole size (Webb and Joss 1997). Interestingly, Pyke and White (2000) also found that the availability of some cover (rocks) did not protect Green and Golden Bell Frogs (Litoria aurea) from attack. Similarly, Baber and Babbit (2004) found that even in the presence of abundant cover, Gambusia were very effective at eating tadpoles of Hyla squirella and Gastrophryne carolinensis, both species which have an evolutionary history with Gambusia. Although I did not feed the Gambusia in my experiment regularly, they lived at relatively low densities and had access to the aquatic invertebrates within the tanks. Moreover, the presence of *Gambusia* in the tanks corresponded with winter, when the metabolic rate (and nutritional requirements) of the fish were minimal. Nevertheless, my results echo those of Baber and Babbit (2004) in demonstrating that the efficiency of Gambusia as predators of anuran larvae is not limited to situations in which Gambusia are new to the community.

This is not to say that *G. holbrooki* has not acted as a selective pressure on *P. feriarum*. For example, adult *P. feriarum* tend not to breed in permanent bodies of water (Skelly 1996), where the danger from predators is highest. *Pseudacris feriarum* tadpoles also alter their behavior in the presence of predators (Skelly 1996). For example, Bridges (2002) showed that *P. feriarum* swam less and used refuges more in the presence of Eastern Newts (*Notopthalmus viridescens*). However, despite a long period of coexistence in southeastern North America, any defenses that *P. feriarum* has evolved

TABLE 1. Relative survival of Pseudacris feriarum in 3	378-1 cattle tanks with and without Gambusia holbrooki.
--	---

Tank Set	Tank	Initial Contents (20 Feb)	Tadpoles Observed (5 Mar)	_
1	А	20 fish, 40 tadpoles	0	_
	В	40 tadpoles	4	
2	А	40 tadpoles	2	
	В	20 fish, 40 tadpoles	0	
3	А	20 fish, 40 tadpoles	0	
	В	40 tadpoles	5	
4	А	40 tadpoles	5	
	В	20 fish, 40 tadpoles	0	
5 A B	А	20 fish, 40 tadpoles	0	
	В	40 tadpoles	6	
6 A B	А	40 tadpoles	2	
	В	20 fish, 40 tadpoles	0	

specifically against *G. holbrooki* appear not to be particularly effective, at least at the early tadpole stage.

Acknowledgments.—I thank John Wear and Catawba College for access to *Gambusia*, Fred Stanback for access to *Pseudacris* eggs, and Michael Dorcas and Malcolm McCallum for comments on the manuscript. The Davidson College IACUC approved this research (protocol number 2-05-12).

LITERATURE CITED

- Baber, M.J., and K. Babbit. 2004. Influence of habitat complexity on predator-prey interactions between the fish (*Gambusia holbrooki*) and tadpoles of *Hyla squirella* and *Gastrophryne carolinensis*. Copeia 2004:173–177.
- Barrier, R.F.G., and B.J. Hicks. 2006. Behavioral interactions between Black Mudfish (*Neochanna diversus* Stokell, 1949: Galaxiidae) and Mosquitofish (*Gambusia affinis* Baird & Girard, 1854). Ecology of Freshwater Fish 3:93–99.
- Bence, J.R. 1988. Indirect effects and biological control of mosquitoes by mosquitofish. Journal of Applied Ecology 25:505–522.
- Bradford, D.F., F. Tabatabai, and D.M. Graber. 1993. Isolation of remaining populations of the native frog, *Rana muscosa*, by introduced fishes in Sequoia and Kings Canyon National Parks, California. Conservation Biology 7:882–888.
- Bridges, C.M. 2002. Tadpoles balance foraging and predator avoidance: Effects of predation, pond drying, and hunger. Journal of Herpetology 36:627–634.
- Brönmark, C., and P. Edenhamn. 1994. Does the presence of fish affect the distribution of tree frogs (*Hyla arborea*)? Conservation Biology 8:841–845.
- Fisher, R.N., and H.B. Shaffer. 1996. The decline of amphibians in California's Great Central Valley. Conservation Biology 10:1387–1397.
- Gamradt, S.C., and L.B. Kats. 1996. Effect of introduced crayfish and mosquitofish on California Newts. Conservation Biology 10:1155–1162.
- Goodsell, J.A., and L.B. Kats. 1999. Effect of introduced Mosquitofish on Pacific Treefrogs and the role of alternative prey. Conservation Biology 13:921–924.
- Grubb, J.C. 1972. Differential predation by *Gambusia affinis* on the eggs of seven species of anuran amphibians. American Midland Naturalist 88:102–108.
- Hamer, A.J., S.J. Lane, and M.J. Mahony. 2002. The role of introduced Mosquitofish (*Gambusia holbrooki*) in excluding the native Green and Golden Bell Frog (*Litoria aurea*) from original habitats in south-eastern Australia. Oecologia 132:445–452.
- Hecnar, S.J., and R.T. M'Closkey. 1997. The effects of predatory fish on amphibian species richness and

distribution. Biological Conservation 79:123-131.

- Kats, L.B., and R.P. Ferrer 2003. Alien predators and amphibian declines: Review of two decades of science and the transition to conservation. Diversity and Distributions 9:99–110.
- Kiesecker, J.M. 2003. Invasive species as a global problem: Toward understanding the worldwide decline of amphibians. Pp. 113–126 *In* Amphibian Conservation. Semlitsh, R. (Ed.). Smithsonian Press, Washington, D.C., USA.
- Knapp, R.A., and K.R. Matthews. 2000. Non-native fish introductions and the decline of the Mountain Yellow-legged Frog from within protected areas. Conservation Biology 14:428–438.
- Komak, S., and M.R. Crossland. 2000. An assessment of the introduced Mosquitofish (*Gambusia affinis holbrooki*) as a predator of eggs, hatchlings and tadpoles of native and non-native anurans. Wildlife Research 27:185–189.
- Kruse, K.C., and M.G. Francis. 1977. A predation deterrent in larvae of the Bullfrog, *Rana catesbeiana*. Transactions of the American Fisheries Society 106:248–252.
- Lawler, S.P., D. Dritz, T. Strange, and M. Holyoak. 1999. Effects of introduced Mosquitofish and Bullfrogs on the threatened California Red-legged Frog. Conservation Biology 13:613–622.
- McCallum, M.L., and S.E. Trauth. 2001. Are tadpoles of the Illinois Chorus Frog (*Pseudacris streckeri illinoensis*) cannibalistic? Transactions of the Illinois State Academy of Science 94:171–178.
- McDiarmid, R.W., and R. Altig. 1999. Tadpoles: The Biology of Anuran Larvae. University of Chicago Press, Chicago, Illinois. USA.
- Miura, T., R.M. Takahashi, and R.J. Stewart. 1979. Habitat and food selection by the Mosquitofish, *Gambusia affinis*. Proceedings of the California Mosquito and Vector Control Association 47:46–50.
- Morgan, L.A., and W.A. Buttermer. 1996. Predation by the non-native fish *Gambusia holbrooki* on small *Litoria aurea* and *L. dentata* tadpoles. Australian Zoologist 30:143–149.
- Moriarty, E., and M.J. Lannoo. 2005. *Pseudacris triseriata* complex. Pp. 485–488 *In* Amphibian Declines: The Conservation Status of United States Species. Lannoo, M. (Ed.). University of California Press, Berkeley, California, USA.
- Petranka, J.W. 1983. Fish predation: A factor affecting the spatial distribution of a stream-breeding salamander. Copeia 1983:624–628.
- Pyke, H.G., and W.W. White. 2000. Factors influencing predation on eggs and tadpoles of the endangered Green and Golden Bell Frog *Litoria aurea* by the introduced Plague Minnow *Gambusia holbrooki*. Australian Zoologist 31:496–505.
- Skelly, D.K. 1996. Pond drying, predators, and the distribution of *Pseudacris* tadpoles. Copeia 1996:599–

Stanback.—Gambusia predation on tadpoles.

605.

- Stebbins, R.C., and N.W. Cohen. 1995. A Natural History of Amphibians. Princeton University Press, Princeton, New Jersey, USA.
- Webb, C., and J. Joss. 1997. Does predation by the fish *Gambusia holbrooki* (Atheriniformes: Pociliidae) contribute to declining frog populations? Australian Zoologist 30:316–324.



MARK STANBACK is a Professor of Biology at Davidson College in Davidson, North Carolina, USA. He received a B.Sc. from Davidson College and a Ph.D. from the University of California at Berkeley. He is an avian behavioral ecologist and is currently studying reproductive strategies in Eastern Bluebirds (*Sialia sialis*), Tree Swallows (*Tachycineta bicolor*), Purple Martins (*Progne subis*), Barn Swallows (*Hirundo rustica*), and Carolina Chickadees (*Poecile carolinensis*). His newest project focuses on conservation of Brown-headed Nuthatches. In addition to teaching Animal Behavior and Evolutionary Biology, he also teaches a Vertebrate Field Zoology course in which herps receive the most attention. (Photographed by Anonymous)