

## AN ASSESSMENT OF TWO METHODS FOR SAMPLING RIVER TURTLE ASSEMBLAGES

SEAN C. STERRETT<sup>1,2,3</sup>, LORA L. SMITH<sup>1</sup>, SARA H. SCHWEITZER<sup>2</sup>, AND JOHN C. MAERZ<sup>2</sup>

<sup>1</sup>Joseph W. Jones Ecological Research Center, Newton, Georgia 39870, USA

<sup>2</sup>Warnell School of Forestry and Natural Resources, University of Georgia, Athens, Georgia 30602, USA

<sup>3</sup>Corresponding author; e-mail: sterretts@warnell.uga.edu

**Abstract.**—We compared the effectiveness of baited hoop traps and effort-managed snorkel surveys for detecting freshwater turtle species within two streams in southwestern Georgia, USA. Snorkel surveys provided the highest detection probability for the four most frequently captured species. We captured the Barbour's Map Turtle (*Graptemys barbouri*), a threatened species in the southeastern U.S., 90% of the time by snorkeling; whereas, we captured the Yellow-bellied Slider (*Trachemys scripta*) 88% of the time by hoop traps. We captured adults and juveniles with both methods, with the exception of juvenile Loggerhead Musk Turtles (*Sternotherus minor*), which we caught only via snorkeling. The two methods captured turtles of similar sizes. Seasonal and physical constraints can affect visibility and access to capture locations, limiting snorkeling sessions. We recommend use of both techniques to capture a range of sexes, sizes, and species of turtles in Coastal Plain streams. However, species-specific surveys may benefit by lower effort with the use of only one method.

**Key Words.**—Coastal Plain; detection probability; Georgia; *Graptemys barbouri*; hoop trapping; snorkel surveys; *Sternotherus minor*, *Trachemys scripta*

### INTRODUCTION

Herpetofauna are notoriously cryptic in habit, yet until recently few studies have accounted for detection probability during research and monitoring efforts. Incorporating measures of detection is essential to making accurate inferences about abundance or a population's response to environmental impacts from comparative or long-term data (MacKenzie and Kendall 2002; Mazzerolle et al. 2007). When working with assemblages of species that differ in behavior, it is unlikely that a single technique will be effective at detecting all species, thus a suite of capture techniques may be required (Liner 2006). Further, various sampling methods can yield differing detection probabilities for any single species (Gunzburger 2007). Nevertheless, researchers are often limited in the numbers of techniques they can employ, so assessments of the effectiveness of individual methods are valuable (Plummer 1979).

Baited hoop traps are commonly used to sample omnivorous, active-foraging freshwater turtles (Legler 1960). However, these methods are generally ineffective for herbivorous and molluscivorous turtles species like *Pseudemys* spp. and *Graptemys* spp., as these species do not respond to bait (MacCulloch and Gordon 1978). Further, hoop traps or fyke nets may be ineffective at capturing small size classes of turtles (Ream and Ream 1966). Alternative methods such as basking traps (Ream and Ream 1966; MacCulloch and Gordon 1978) and fyke nets (Vogt 1980) are used to capture species that do

not readily come to bait (Plummer 1979; Browne and Hecnar 2005). Not surprisingly, studies of whole turtle assemblages rely on multiple sampling methods, although the detection probabilities of the individual and combinations of techniques are seldom reported (Dreslik et al. 2005; Smith et al. 2006; Browne and Hecnar 2007).

Hand capture via snorkeling ("goggling") has been used to supplement other methods of aquatic turtle capture (Marchand 1945; Chaney and Smith 1950; Allen and Neil 1950). This method is particularly effective in clear, shallow lakes and rivers. For example, Carr (1952) recounts a goggling trip of Lewis Marchand and colleagues in which 60 Barbour's Map Turtles (*Graptemys barbouri*) were captured over three days on the Chipola River in northwest Florida, and Marchand (1945) captured 163 *Pseudemys* spp. via snorkeling in Rainbow Run, a spring-fed river in west-central Florida. Polisar (1995) found that snorkeling for *Dermatemys mawii* resulted in more captures than any other method used in tributaries of the Belize River. Further, snorkeling yielded captures of all size classes of *D. mawii*. Though snorkeling may provide greater detection of some turtle species and size classes, the technique can be limited by dynamic water conditions and the difficulty in standardizing captures across observers with different abilities. For example, Polisar (1995) described several challenges with free diving for turtles, such as seasonal fluctuations in visibility and stream depth and the variation in the observer's ability to reach all depths to search for turtles.

The objectives of the present study were to compare species detection rates and assess capture by species and sex for snorkeling and baited hoop trapping for an assemblage of freshwater turtle species in southwestern Georgia. Streams in the southeastern United States, particularly those that drain into the Gulf of Mexico, are among the richest in turtle diversity in the world (Buhlmann and Gibbons 1997). Streams in southwestern Georgia contain as many as nine turtle species (Jensen et al. 2008). Georgia's State Wildlife Action Plan has identified monitoring populations and determining the effects of human activities on the composition and abundance of freshwater turtles as a priority conservation action (Georgia Department of Natural Resources, <http://www.gadnr.org/cwcs/Documents/strategy.html>). These objectives can be met with more rigor when capture techniques have been thoroughly assessed.

## MATERIALS AND METHODS

**Study area.**—This study took place on Ichawaynochaway (Baker County, Georgia) and Spring Creeks (Decatur and Miller counties, Georgia) in the Lower Flint River Basin (LFRB) of southwestern Georgia. The creeks are located in the Dougherty Plain physiographic district, which is characterized by karst topography (Ward et al. 2005). Southwestern Georgia has a high variability in annual rainfall with an average of 1270 mm per year and precipitation and stream flows are lowest during summer (Golden and Hess 1991; Golladay and Battle 2002). In drainages of the LFRB, rocky limestone shoals and deep, wide, sandy pools are common. Both creeks have ground water input from the Upper Floridan Aquifer (Hicks, D.W., and S.W. Golladay. 2006. Impacts of agricultural pumping on selected streams in southwestern Georgia. Submitted to Georgia Environmental Protection Division, Atlanta, Georgia). During late spring and summer, both creeks are clear enough to see the bottom in most areas unless there have been recent rain events, which reduce visibility.

**Turtle surveys.**—We randomly selected seven 1-km study segments on each creek (14 total sites) and a 0.5-km section in the center of each 1-km segment as the focus of survey effort. We surveyed from June to August 2007 on Ichawaynochaway Creek and June to September 2008 on Spring Creek. We trapped using baited hoop traps over five consecutive nights twice during the sampling period and used snorkel surveys twice over the sampling period at each study section (for more detail, see Sterrett et al. in press). In most cases, both sampling methods were independent although due to water clarity limitations, these methods did overlap on several occasions.

During each survey, we placed five large (4-hoops 1.2-m diameter, 3.8-cm mesh size) and five small (3-hoops 0.9-m diameter, 3.8-cm mesh) hoop traps (Memphis Net and Twine, Memphis, Tennessee, USA) baited with fish in the center 0.5-km section of each 1-km study segment. We set traps approximately 50 m apart on alternating banks in locations where traps were mostly inundated by water. Traps were run for five consecutive nights and we then checked daily and rebaited as necessary.


During snorkel surveys, we standardized search effort as follows: 3–4 surveyors for 2–3 h (actual time was recorded), time of day (1300 EST start time, when possible), and surveyor experience, with 1–2 experienced surveyors and 1–2 volunteer surveyors on each survey. At least one experienced surveyor with extensive experience snorkeling and capturing turtles was present for all surveys. Volunteers were all wildlife biologists that varied in experience with freshwater turtles and snorkeling and contributed only a few days each to sampling. We quantified capture success as the number of turtles captured per person-hour. During each survey, we searched the study section twice (once upstream and once downstream). Surveyors thoroughly searched all accessible areas within each stream section. Particularly high numbers of captures in one section limited sampling in that section to one upstream search.

We measured maximum straight-line carapace length (CL), maximum plastron length (to the nearest 1 mm), and body mass (to the nearest 10 g) for each captured turtle. We gave each turtle a unique identification code by marking the marginal scutes (Cagle 1939), except softshell turtles (*Apalone* spp.), which were marked with zip-ties or notches following Plummer (2008).

**Data analyses.**—We used program PRESENCE (MacKenzie and Kendall 2002; Hines 2006) to estimate the single season detection probability ( $p$ ) for the four most frequently captured turtle species for each method, each year. To use PRESENCE, at least 60 captures per species were required, so we did not calculate detection probabilities for infrequently captured species. In the analysis, we pooled data from both creeks and calculated sampling occasions as trap-nights [ $N = (10 \text{ traps set for } 5 \text{ nights})(2 \text{ trapping sessions})(7 \text{ replicates } [0.5\text{-km sections}])(2 \text{ creeks}) = 1400$ ] and snorkeling visits [ $N = (2 \text{ snorkeling sessions})(7 \text{ replicates } [0.5\text{-km sections}])(2 \text{ creeks}) = 28$ ]. To calculate detection probability for both methods combined, we used the equation,

$$p_{\text{both methods}} = 1 - (1 - p_{\text{method 1}})(1 - p_{\text{method 2}})$$

(described in Farmer et al. 2009), where  $p_{\text{method 1}}$  and  $p_{\text{method 2}}$  were obtained from program PRESENCE. Standard error for detection probability estimates for



**FIGURE 1.** Mean detection probabilities ( $\pm$ SE) of the four most frequently captured turtle species by sampling method and for both methods combined. An uncapped standard error bar indicates a standard error of  $>1$ . Sampling was conducted on Ichawaynochaway (Baker County) and Spring Creeks (Decatur and Miller counties) in the lower Flint River Basin of southwestern Georgia, USA, in 2007–2008.

each species and each method were calculated using the delta method of approximating standard error (Williams et al. 2002). We used Chi-square analysis to determine if the sex ratio of captured turtles differed among species. Means were considered significantly different at  $\alpha = 0.10$ . We used a general linear model to investigate if capture method, sex, or species (independent variables) had any influence on carapace length of adults (dependent variable). We did not differentiate turtle capture by trap type until late in the study, so we could not examine the differences in detection probability between large and small hoop traps.


## RESULTS

We trapped for 10 nights per study section (1400 trap nights total), which required approximately 10–20 person hours to set and check traps at each site. Effort-managed snorkeling required two 3-hour visits (9–12 person hours) per site (242.75 total person hours). We captured 674 individual turtles (823 total captures) representing nine species on the two creeks over the course of the study. Four species (Yellow-Bellied Slider, *Trachemys scripta*; Barbour's Map Turtle, *Graptemys barbouri*; River Cooter, *Pseudemys concinna*; Loggerhead Musk Turtle, *Sternotherus minor*) comprised 95% of all individual captures. We captured at least one individual of all nine species by both methods. We captured most (90%,  $n = 102$ ) *G. barbouri* by snorkeling, whereas we captured most (88%,  $n = 301$ ) *T. scripta* in traps. We captured 20 *T. scripta* and two *G. barbouri* with one method and later recaptured them

with another method during the study. Captures by traps and snorkel surveys were similar for *P. concinna* (40% and 60%, respectively,  $n = 120$ ) and *S. minor* (53% and 47%,  $n = 70$ ). Ten of 14 recaptured *P. concinna* were captured with both methods. We only caught one *S. minor* with both methods. We caught most Alligator Snapping Turtles (*Macrochelys temminckii*; 72%,  $n = 18$ ) by trapping. We caught Spiny Softshell Turtles (*Apalone spinifera*;  $n = 14$ ), Florida Softshell Turtles (*A. ferox*;  $n = 1$ ), and Common Snapping Turtles (*Chelydra serpentina*;  $n = 3$ ) only in traps. All Florida Cooters (*P. floridana*;  $n = 7$ ) were captured by snorkeling.


Detection probabilities for the four most frequently captured species were all greater for effort-managed snorkeling than for baited hoop trapping (Fig. 1). *Graptemys barbouri*, *P. concinna*, and *S. minor* all had similarly low detection probabilities with baited hoop trapping (Fig. 1). Baited hoop trapping produced a similar detection probability for *T. scripta* compared to snorkeling. The combination of snorkeling and baited hoop trapping did not significantly improve detection compared to snorkeling alone for any of the four most frequently captured species (Fig. 1).

With one exception, the distributions of male, female and juvenile turtles caught by snorkeling and baited hoop trapping were similar for the four most common turtle species (snorkeling  $\chi^2 = 43.3$ ,  $df = 6$ ,  $P = 0.00$ ; trapping  $\chi^2 = 16.9$ ,  $df = 6$ ,  $P = 0.01$ ; Fig. 2). The exception was that juveniles of *S. minor* were detected using snorkeling but were never captured in a baited hoop trap. More individuals of each sex of *T. scripta* were captured by trapping, while more individuals of each sex of *G. barbouri* were captured by snorkeling



**FIGURE 2.** Number of turtles captured by sex for the four most frequently captured species on Ichawaynochaway and Spring Creeks, Georgia, USA, in 2007–2008. Sampling methods included A.) Large and small hoop traps ( $\chi^2 = 16.9$ ,  $df = 6$ ,  $P = 0.01$ ) and B.) Effort-managed snorkeling ( $\chi^2 = 43.3$ ,  $df = 6$ ,  $P = 0.00$ ).

(Fig. 2). The overall model showed that species and sex had significant effects on CL ( $F = 243.9$ ,  $df = 3$ ,  $P = 0.00$  for species;  $df = 1$ ,  $F = 152.86$ ,  $P = 0.00$  for sex; Fig. 3). However, method did not have a significant effect on CL of the four most frequently captured species ( $F = 0.021$ ,  $df = 1$ ,  $P = 0.88$ ; Fig. 3).



**FIGURE 3.** Mean carapace length of the four most frequently captured turtles by hoop trapping and effort-managed snorkeling on Ichawaynochaway and Spring Creeks, Georgia, USA, in 2007–2008. Error bars represent standard error.

## DISCUSSION

The baited hoop net is a standard trap for freshwater turtles and is considered a reliable method for detecting many species (Plummer 1979). In our study, across all sampled habitat (7 km of creek), baited hoop trapping or snorkeling were both sufficient to detect all nine species of turtles expected to be present in the study streams; however, snorkel surveys yielded higher detection rates for the four most common species at the finer scale (within a 0.5 km section) and yielded captures of all obligate riverine species with the exception of *A. spinifera*. Baited hoop trapping yielded a detection probability similar to that of snorkel surveys only for *T. scripta* and the combination of the two methods had no measurable increase in detection over snorkeling alone. Therefore, our results suggest that baited hoop trapping is inferior to snorkeling for detecting river turtles in this system, particularly for rare species or species with habits that do not draw them to bait. When considering the effectiveness of a method, it is important to take into account the time, effort, and expertise required relative to the detection probability achieved. Further, the choice of method used in a particular system should be species specific.

With one exception, the two methods did not differ in the distributions of sexes or size classes of turtles. The one exception was the failure to capture juvenile *S. minor* in traps. This was likely due to their small size (hatchlings are 22–30 mm in CL; Ernst and Lovich 2009) and ability to slip through the mesh size of the standard hoop traps we used. Despite perceptions that small turtles are more cryptic than large turtles (Carr

1952), we found that small turtles were easily seen at the base of logs and twigs when snorkeling.

While our results suggest that snorkeling yields higher detection probabilities than baited hoop trapping with less effort (two days versus five), snorkeling may not be as effective in other contexts. Ichawaynochaway Creek and Spring Creek are relatively clear water bodies with excellent visibility on most summer days, when precipitation and flow are low. More degraded water bodies, which are arguably more common, are likely to have poor visibility, limiting the effectiveness of snorkeling. In these situations, baited hoop trapping may still be the most effective method for sampling turtles. Because detection can be low and vary among species, use of baited hoop traps should account for detection probability in predicting habitat occupancy by turtle species (*sensu* Rizkalla and Swihart 2006). In situations where baited hoop traps are the best option, methods such as basking traps or basking surveys should supplement trapping to detect species not attracted by bait.

Further, while the focus of this analysis was on detection of species within a section of creek, we acknowledge that the most efficient methods may differ if the objective is to capture large numbers of individuals for population estimation. Koper and Brooks (1998) report that neither baited hoop traps, basking surveys, nor hand captures were adequate for estimating turtle population size, but that a combination of methods improved the accuracy of estimates. Multiple methods tend to yield greater numbers of captures than any single method. In our study, neither of the two methods alone maximized captures for all species. Baited hoop traps captured many more *T. scripta* of both sexes, indicating that this method would be most useful in estimation of *T. scripta* population size. By contrast, snorkeling captured more *G. barbouri* of both sexes, indicating that this technique would be better suited to accurately estimating the species' abundance. Few *M. temminckii* ( $n = 18$ ) were captured during the course of the study, suggesting that either the combined effort of both methods was not robust enough to capture large numbers of *M. temminckii* or densities are relatively low in this area (Jensen and Birkhead 2003). Greater sampling effort (more traps and/or more snorkeling surveys) might be required to capture sufficient numbers of this species to estimate population size. The surprisingly few captures of *A. spinifera* during the study remain a mystery. While conducting snorkel surveys, we observed two adult *A. spinifera*, but were unable to capture either individual. *Apalone spinifera* are known for their rapid swimming, which made it challenging for surveyors to make a capture once they had observed an individual. However, this species is often captured readily in baited hoop traps (Plummer 1979). It is possible that this species also

occurs in low densities in the creeks sampled and that more trapping effort would be required to yield captures.

We did not attempt to use three other common methods for surveying turtles in this study: basking traps, direct observation of basking turtles, and fyke nets. Koper and Brooks (1998) suggested that basking observations are an efficient and unbiased way to estimate population size for *Chrysemys picta*. Basking traps and fyke nets are also effective for turtles not attracted to bait (Plummer 1979). We feel that at the scale of our study, the effort needed to use basking traps would not have improved detection and may not have increased greatly the numbers of turtles observed. Setting up basking traps and monitoring them would require a similar or greater effort than baited hoop trapping, and a greater effort than snorkeling. Further, basking trap effort would be difficult to standardize because of the need for available basking substrates to attach traps, and would be biased against species that bask under the surface of the water. We also did not attempt to install fyke nets in the stream sections because of the inability to reliably set these in limestone substrate and because of occasional high flows which may have damaged or dislodged nets during times of sampling. Vogt (1980) found fyke nets highly effective in pond and backwater areas, but noted their inconvenience in faster current.

Our results show that the most common method used in studies of aquatic turtles, baited hoop trapping, can have lower power to detect some species at finer spatial scales. We found that snorkel surveys provided a better level of detection than hoop trapping for all turtle species; however, hoop trapping did yield higher numbers of *T. scripta*. Collectively, the results suggest that the most effective method for sampling freshwater turtles will depend on the focal species and study objectives. Multiple capture methods may be required if the objectives are to accurately measure both habitat occupancy and abundance. Further, the most effective method is likely to vary depending on whether local conditions limit opportunities to use methods such as snorkel surveys.

*Acknowledgments.*—The funding for this project was made possible by a Georgia Department of Natural Resources State Wildlife Grant in coordination with the United States Fish and Wildlife Service and matching funds from the Joseph W. Jones Ecological Research Center. We thank Brent Howze, David Steen, Tara Muenz, Andrew Grosse, Gabriel Miller, Aubrey Heupel, Carla Atkinson, Jen Linehan, Kelley McKean, Steven Jones, Anna Derrick, Greg Knothe, Lindsey Sargent, Evan Hill, Amy Krzton-Presson, Stephen Golladay, Scott Stavely, Michael Mazzacavallo, and Jessica Gonynor for assistance in snorkeling. Liz Cox assisted in acquiring literature. Thanks to David A. Steen and

Kyle Barrett for reviewing earlier drafts of this manuscript. All procedures in this study were approved by the Georgia Animal Care and Use Committee (AUP # A2007-10102-0) and under the Georgia Department of Natural Resources, Scientific Collecting Permit, 29-WCH-06-104.

## LITERATURE CITED

- Allen, E.R., and W.T. Neill. 1950. The Alligator Snapping Turtle, *Macrochelys temminckii*, in Florida. Special Publication of Ross Allen's Reptile Institute 4:1–15.
- Browne, C.L., and S.J. Hecnar. 2005. Capture success of Northern Map Turtles (*Graptemys geographica*) and other turtle species in basking vs. baited hoop traps. *Herpetological Review* 36:145–147.
- Browne, C.L., and S.J. Hecnar. 2007. Species loss and shifting population structure of freshwater turtles despite habitat protection. *Biological Conservation* 138:421–429.
- Buhlmann, K.A., and J.W. Gibbons. 1997. Imperiled Aquatic Reptiles of the Southeastern United States: Historical Review and Current Conservation Status. Pp. 201–231 *In* *Aquatic Fauna in Peril: The Southeastern Perspective*. Benz, G.W., and D.E. Collins (Eds.). Southeast Aquatic Research Institute, Decatur, Georgia, USA.
- Cagle, F.R. 1939. A system of marking turtles for future identification. *Copeia* 1939:170–172.
- Carr, A. 1952. *Handbook of Turtles: Turtles of the United States, Canada, and Baja California*. Comstock Publishing Association, Ithaca, New York, USA.
- Chaney, A., and C.L. Smith. 1950. Methods for collecting map turtles. *Copeia* 1950:323–324.
- Dreslik, M.J., A.R. Kuhns, and C.A. Phillips. 2005. Structure and composition of a southern Illinois freshwater turtle assemblage. *Northeastern Naturalist* 12:173–186.
- Ernst, C.H., and J.E. Lovich. 2009. *Turtles of the United States and Canada*. The Johns Hopkins University Press, Washington, D.C, USA.
- Farmer, A. L., L. L. Smith, S. B. Castleberry, and J. W. Gibbons. 2009. A comparison of techniques for sampling amphibians in isolated wetlands in Georgia, USA. *Applied Herpetology* 6:327–341.
- Golden, H.G., and W.G. Hess. 1991. Georgia: Floods and Droughts. Pp 239–243 *In* *National Water Survey 1988–1999*. U.S. Geologic Survey, Water-Supply Paper 2375, USGS, Reston, Virginia, USA.
- Golladay, S.W., and J. Battle. 2002. Effects of flooding and drought on water quality in Gulf Coastal Plain Streams in Georgia. *Journal of Environmental Quality* 31:1266–1272.
- Gunzburger, M.S. 2007. Evaluation of seven aquatic sampling methods for amphibians and other aquatic fauna. *Applied Herpetology* 4:47–63.
- Hines, J.E. 2006. PRESENCE2 – Software to estimate patch occupancy and related parameters. USGS-PWRC. Available from <http://www.mbr-pwrc.usgs.gov/software/-presence.html>.
- Jensen, J.B., and W.S. Birkhead. 2003. Distribution and status of the Alligator Snapping Turtle (*Macrochelys temminckii*) in Georgia. *Southeastern Naturalist* 2:25–34.
- Jensen, J.B., C.D. Camp, J.W. Gibbons, and M.J. Elliott. 2008. *Amphibians and Reptiles of Georgia*. University of Georgia Press, Athens, Georgia, USA.
- Koper, N., and R.J. Brooks. 1998. Population-size estimators and unequal catchability in painted turtles. *Canadian Journal of Zoology* 76:458–465.
- Legler, J.M. 1960. A simple and inexpensive device for trapping aquatic turtles. *Proceedings of the Utah Academy of Sciences, Arts and Letters* 37:63–66.
- Liner, A.E. 2006. Wetland predictors of amphibian distributions and diversity within the Southeastern U.S. Coastal Plain. M.S. thesis, University of Georgia, Athens, Georgia, U.S.A. 112 p.
- MacCulloch, R.D., and D.M. Gordon. 1978. A simple trap for basking turtles. *Herpetological Review* 9:133.
- Marchand, L.J. 1945. Water goggling: a new method for the study of turtles. *Copeia* 1945:37–40.
- Mazzerolle, M.J., L.L. Bailey, W.L. Kendall, J.A. Royle, S.J. Converse, and J.D. Nichols. 2007. Making great leaps forward: accounting for detectability in herpetological field studies. *Journal of Herpetology* 41:672–689.
- MacKenzie, D.I., and W.L. Kendall. 2002. How should detection probability be incorporated into estimates of relative abundance? *Ecology* 83:2387–2393.
- Plummer, M.V. 1979. Collecting and Marking. Pp. 45–60 *In* *Turtles: Perspectives and Research*. Harless, M., and H. Morlock (Eds.). John Wiley & Sons, Inc., New York, New York, USA.
- Plummer, M.V. 2008. A notching system for marking softshell turtles. *Herpetological Review* 39:64–65.
- Polisar, J. 1995. River turtle reproductive demography and exploitation patterns in Belize: implications for management. *Vida Silvestre Neotropical* 4:10–19.
- Ream, C., and R. Ream. 1966. Influence of sampling methods on estimation of population structure in painted turtles. *American Midland Naturalist* 75:325–338.
- Rizkalla, C.E., and R.K. Swihart. 2006. Community structure and differential responses of aquatic turtles to agriculturally induced habitat fragmentation. *Landscape Ecology* 21:1361–1375.
- Smith, G.R., J.B. Iverson, and J.E. Rettig. 2006. Changes in a turtle community from a northern Indiana lake: A long-term study. *Journal of Herpetology* 40:180–185.

Sterrett et al.—Assessment of river turtle sampling methods.

- Sterrett, S.C., L.L. Smith, S.W. Golladay, S.H. Schweitzer, and J.C. Maerz. In press. The conservation implications of riparian land use on river turtles. *Animal Conservation*.
- Vogt, R.C. 1980. New methods for trapping aquatic turtles. *Copeia* 1980:368–371.
- Ward, G.M., P.M. Harris, and A.K. Ward. 2005. Gulf Coast Rivers of the Southeastern United States. Pp. 125–178 *In* Rivers of North America. Benke, A.C., and C.E. Cushing (Eds.). Elsevier Academic Press, San Diego, California, USA.
- Williams, B.K., J.D. Nichols, and M.J. Conroy. 2002. Analysis and Management of Animal Populations: Modeling, Estimation and Decision Making. Academic Press, San Diego, California, USA.

## Herpetological Conservation and Biology



**SEÁN C. STERRETT** is a Ph.D. student at the Warnell School of Forestry and Natural Resources at the University of Georgia. He began studying turtles at Butler University, where he received a Bachelors of Science in Biology in 2003 and studied urban populations of freshwater turtles. Sean migrated to the Joseph W. Jones Ecological Research Center where he worked as a Herpetology Research Technician, then completed his Masters degree in 2009 in conjunction with the Warnell School, studying the effects of land use on river turtle assemblages. His current research interests include freshwater turtle ecology and conservation, with an emphasis on map turtles. Sean is also co-president of the Society of Conservation Biology Georgia Chapter. (Photographed by Aubrey M. Heupel)



**LORA L. SMITH** is an Associate Scientist at the Joseph W. Jones Ecological Research Center in southwestern Georgia. Her work at the Center includes a long-term study of the effects of mammalian predators on Gopher Tortoises (*Gopherus polyphemus*), upland snake ecology, and habitat predictors of pond-breeding amphibians. Her research interests also include the effects of habitat restoration and management on amphibian and reptiles. (Photographed by Kelly McKean)



**SARA W. SCHWEITZER** completed a B.S. in Biology from the University of North Carolina, Chapel Hill; a M.S. in Wildlife Science from Texas Tech University, Lubbock; and a Ph.D. in Wildlife Ecology from Oklahoma State University, Stillwater. Between her M.S. and Ph.D., she worked as an Environmental Scientist for the U.S. Army Corps of Engineers, Regulatory and Waterways Sections, Norfolk, Virginia. Currently, Dr. Schweitzer is a Professor within the Warnell School of Forestry and Natural Resources, University of Georgia, Athens. Her research focuses on responses of wildlife populations and habitats to anthropogenic disturbances, especially within wetland and coastal areas of the South Atlantic Coastal Plain. Her current research projects include wintering ecology of Rusty Blackbirds (*Euphagus carolinus*), wintering and nesting ecology of King Rails (*Rallus elegans*) and Clapper Rails (*Rallus longirostris*), and adaptations of Wood Storks (*Mycteria Americana*) nesting within an urban environment. She also has contracts with the USFWS and US Department of Defense to survey and monitor for rare species on military lands, including the Gopher Tortoise and Indigo Snake (*Drymarchon couperi*). (Photographed by John Seginak)



**JOHN C. MAERZ** is an Associate Professor of vertebrate ecology in the Warnell School of Forestry and Natural Resources at the University of Georgia. He came to UGA in 2005 after four years as a research associate in the Department of Natural Resources at Cornell University. He received a B.S. in Zoology from the University of Maryland and a Ph.D. in Biology from the State University of New York at Binghamton. Dr. Maerz did his postdoctoral work in the Ecology and Management of Invasive Plants Program at Cornell University where he started several research projects examining the impacts of non-native plant and invertebrate invasions on amphibians in terrestrial and freshwater ecosystems. At the University of Georgia, his research program uses amphibians and reptiles to understand the effects of terrestrial and aquatic environmental change on wildlife. He continues to focus on the effects of nonnative species invasions, and has broadened his focus to the impacts of agriculture exurban residential development. Dr. Maerz has also started a research program to understand the affects of salamanders and turtles on nutrient dynamics in freshwater ecosystems. (Photographed by Vanessa C. Kinney)