AVOIDANCE OF THREE HERBICIDE FORMULATIONS BY EASTERN RED-BACKED SALAMANDERS (*Plethodon cinereus*)

BENJAMIN J. GERTZOG, LINDSAY J. KAPLAN, DANIEL NICHOLS, GEOFFREY R. SMITH¹, AND JESSICA E. RETTIG

Department of Biology, Denison University, Granville, Ohio 43023 ¹Correspnding author, e-mail: smithg@denison.edu

Abstract.—The ability of terrestrial amphibians to avoid chemical pollutants, such as herbicides or fertilizers, is relatively unknown although such chemicals may occur in their habitats. We studied whether the Eastern Red-backed Salamander (*Plethodon cinereus*) would be able to behaviorally avoid three different herbicide formulations (Spectracide Brush Killer, Bayer Advanced Brushkiller Plus, and Roundup Weed and Grass Killer Concentrate Plus). Salamanders avoided all three formulations when encountered at their full recommended application concentration, and were able to detect and avoid the Bayer and Roundup formulations at 10% the full concentration. Salamanders did not avoid any formulation at 1% the full concentration. Our results suggest that salamanders are able to detect these herbicide formulations and behaviorally avoid them, perhaps reducing any negative effects from exposure to some field applications.

Key Words.---amphibian; avoidance behavior; Eastern Red-backed Salamander; herbicide; Plethodon cinereus

INTRODUCTION

Herbicides are frequently used in forested habitats to help manage timber production (Shepard et al. 2004). In addition, herbicide drift into forests from neighboring agricultural fields can also occur (Gove et al. 2007). Such herbicide use and drift can occur over large areas. However, herbicides also can be used on a smaller scale in natural areas to control invasive plants (Carlson and Gorchov 2004; Simmons et al. 2007; Slaughter et al. 2007). The widespread use of these herbicides has raised questions about their potential effects on nontarget organisms, including both plants and animals (Guynn et al. 2004). In particular, the effects on forestdwelling amphibians, both aquatic and terrestrial, have been a subject of concern.

Unfortunately, we have relatively little information on such effects for terrestrial amphibians. Relyea (2005) found that overspraying of Roundup killed the majority of juvenile anurans tested, including Lithobates sylvaticus, Anaxyrus fowleri, and Hyla versicolor. Dinehart et al. (2009) also found increased mortality in juvenile anurans (Anaxyrus cognatus and Spea multiplicata) exposed to Roundup Weed and Grass Killer Ready-to-Use Plus, and Weed and Grass Killer Super Concentrated, but were not killed by Roundup Weather Max and Ignite 280SL. Agroxone 5 (formulation of 2,4-D) was lethal to adult Triturus cristatus carnifex (Zaffaroni et al. 1986). However, Cole et al. (1997) found that spraying with only glyphosate did not affect capture rates in several terrestrial salamanders, including Plethodon vehiculum and P.

dunni. Glyphosate alone did not affect several species of terrestrial amphibians from the Oregon Coast Range (McComb et al. 2008), but commercial formulations using surfactants were not tested. Similarly, Jones et al. (2000) found little effect of herbicide treatment (tebuthimon) on amphibians and reptiles and Bernal et al. (2009) also found little effect of glyphosate formulations on Colombian frogs at normal application rates. Little is known about the ability of terrestrial amphibians to behaviorally avoid these herbicides as a means to potentially minimize the toxic effects. In one of the few published studies, Storrs Méndez et al. (2009) found that juvenile American Toads (Anaxyrus americanus) do not avoid soils treated with atrazine. However, adult amphibians may be able to detect and avoid herbicides, as evidenced by the ability of ovipositing anurans to avoid depositing eggs in contaminated pools or ponds (e.g., Takahashi 2007; Vonesh and Buck 2007; Vonesh and Kraus 2009).

We examined whether or not Eastern Red-backed Salamanders (*Plethodon cinereus*) would avoid substrates contaminated with three common household herbicide formulations (Spectracide Brush Killer Bayer Advanced Brushkiller Plus and Roundup Weed and Grass Killer Concentrate Plus), and thus whether or not they might be able to avoid exposure to lethal or sublethal concentrations of herbicides in nature if they are able to detect them. Because Eastern Red-backed Salamanders are frequently the most abundant terrestrial vertebrate in the forests of eastern North America (Casper 2005), understanding how they might behaviorally respond to herbicide use is very important, particularly given their important role in many temperate forest ecosystems (Welsh and Droege 2001; Davic and Welsh 2004). In addition, other less common species of *Plethodon* are also likely to experience such stressors and thus an understanding of the response of *P. cinereus* to herbicides will provide some understanding of how these other, more threatened, species may respond.

MATERIALS AND METHODS

We collected adult male and female Eastern Red Backed Salamanders (Plethodon cinereus; SVL about 45-55 mm) during mid-April 2010 and 2011 from under cover boards and natural cover objects on the Denison University Biological Reserve in Granville, Licking Co., Ohio, USA. The three studied herbicides were Spectracide Brush Killer (United Industries Corp., St. Louis, Missouri, USA; 2,4-D, 2-ethylexyl ester 9.74%, Dichlorprop-p, 2-ethylexyl ester 4.78%, Dicamba 1.65%; surfactant unkown), Bayer Advanced Brushkiller Plus (Bayer Crop Science LP, Research Triangle Park, North Carolina, USA; Triclopyr, triethylamine salt 8.8%; surfactant unknown), and Roundup Weed and Grass Killer Concentrate Plus (Monsanto Company, Marvsville, Ohio, USA: Glyphosate, isopropylamine salt 18%, Diquat dibromide 0.73%; surfactant POEA).

We used 15 cm diameter petri dishes as experimental arenas. We lined the bottom of each petri dish with two halves of a 15 cm diameter filter paper. One half was dipped into aged tapwater and the other half into the herbicide formulation. We determined full herbicide concentrations using the recommended dilutions provided by the manufacturer on the label: Bayer (31.25 mL/L), Spectracide (39.1 mL/L), and Round up (46.9 mL/L). We ran separate choice tests on the full, 10%, and 1% concentrations of each herbicide formulation. In addition, we conducted a double control set of trials in which both sides of the filter paper were treated with aged tapwater. We did not determine actual concentrations of the herbicides. For the full concentration and control trials, we used 35 salamanders per formulation (total number of salamanders used = 140). For the 10% and 1% concentration trials, we used 20 salamanders per formulation and concentration (total number of salamanders used = 120). No salamanders were used more than once.

For each trial, we placed a salamander in the middle of the petri dish. When each of the salamanders was in their respective dishes, we covered the dishes and started timers. We recorded the location of the salamander on either the control or herbicide side of the petri dish every 15 min for 1 h. We determined the location of the salamander by where the majority of the front portion of their body lay. For each individual, we calculated the proportion of observations the salamander was on the water side of the petri dish. We used a Wilcoxon

signed-rank test to compare the mean proportion of observations on the water side to the expected mean of 0.5 for random use of the sides of the petri dish. We ran separate tests for each herbicide and concentration. Means are given ± 1 S.E.

RESULTS

For our double control trials, we found that there was no preference for either side of the Petri dish (mean proportion on left side = 0.54 ± 0.08 ; W = 24.0, df = 34, P = 0.60).

At the full concentration of Spectracide, we observed salamanders on the water side of the petri dish significantly more than would be expected by chance (mean = 0.76 ± 0.057 ; W = 186, df = 34, *P* < 0.001). Salamanders did not show a significant preference for water or Spectracide sides of the petri dish at either the 10% (mean = 0.39 ± 0.08 ; W = -34.5, df = 19, *P* = 0.13) or the 1% concentration (mean = 0.46 ± 0.06 ; W = -6.0, df = 19, *P* = 0.75).

In the full concentration trial for the Bayer formulation, salamanders significantly avoided the Bayer-treated side of the petri dish (mean = 0.81 ± 0.057 ; W = 225.5, df = 34, *P* < 0.001). The salamanders also significantly avoided the Bayer-treated side of the petri dish at 10% concentration (mean = 0.70 ± 0.08 ; W = 52.5, df = 19, *P* = 0.049). Salamanders did not show any preference at the 1% concentration (mean = 0.48 ± 0.09 ; W = -4.5, df = 19, *P* = 0.92).

Salamanders significantly avoided the side of the petri dish treated with the full concentration of the Round-up formulation (mean = 0.69 ± 0.07 ; W = 127.5, df = 34, *P* = 0.009). Salamanders also avoided the Round-up treated side of the petri dish at the 10% concentration (mean = 0.76 ± 0.07 ; W = 69.5, df = 19, *P* = 0.003). No preference was observed at the 1% concentration (mean = 0.59 ± 0.07 ; W = 11.5, df = 19, *P* = 0.29). All salamanders survived the experiment.

DISCUSSION

Our results suggest that Eastern Red-backed Salamanders are able to detect and avoid all three herbicide formulations at their full label application rates, and to avoid both the Bayer and Round-up formulations at 10% the label concentration. The actual avoidance response in the field may take the form of dispersal from the area, as has been observed in response to clear-cutting (Semlitsch et al. 2008), or by taking refuge belowground. The ability to avoid these herbicide formulations would potentially allow Redbacked Salamanders to minimize exposure to the potentially toxic effects of these herbicides which have been shown to be lethal to some species of terrestrial amphibians (e.g., Zaffaroni et al. 1986; Relyea 2005; Dinehart et al. 2009). However, the inability of these salamanders to respond to more dilute concentrations may limit their ability to avoid exposures to these herbicides. For example, our results suggest Eastern Red-backed Salamanders could avoid exposure to herbicides from small-scale spot treatment of the understory which would be expected to result in higher concentrations at ground level, but might not be able to avoid exposure to larger scale applications, such as aerial treatments, which would be expected to result in lower concentrations at ground level.

We used commercially-available formulations of the herbicides that contained both active ingredients and surfactants. The response we observed could therefore be driven by either the active ingredients or the surfactants. We suspect that the surfactants are likely the major factor in the salamanders' responses since in several other contexts the surfactants of herbicides have been shown to be more toxic to amphibians than the active ingredients (e.g., Mann and Bidwell 2001; Mann et al. 2003; Howe et al. 2004). Unfortunately, the specific surfactant was known only for one of the formulations we used (Roundup), and thus we cannot generalize about the relative effects of different surfactants and their formulations.

In addition to the results of our study, *P. cinereus* have also been shown to avoid soils contaminated with high levels of lead (Bazar et al. 2010) and urea (Gaglione et al., unpubl. data) and to choose substrates based on pH (Mushinsky and Brodie 1975). Thus, Eastern Redbacked Salamanders appear to have the ability to detect and avoid a wide range of contaminants, which might be expected given the ability of these salamanders to detect and respond to chemical cues from conspecifics and predators (Maerz et al. 2001; Sullivan et al. 2002; Martin et al. 2005; Dantzer and Jaeger 2007).

From a management perspective, our results suggest that Eastern Red-backed Salamanders will be able to behaviorally respond to herbicide treatments that may be used in controlling invasive species or other weeds in forest habitats, especially if such use is targeted and used on relatively small areas. Unfortunately, we know too little about the toxicity of field applications of herbicides to know if temporary avoidance of herbicides would be effective, or even if avoidance behaviors and the salamander's mobility would allow salamanders to escape the toxic herbicides when used on a larger scale. In addition, it is unclear if there may be any negative consequences for the avoidance behavior, such as reduced foraging or increased risk of predation due to greater movement.

Acknowledgments.—We thank Wesley Smith for help collecting the salamanders and running some of the 2011 trials. We also thank Michelle Boone and an anonymous reviewer for their helpful comments on this manuscript.

This experiment was approved by the Denison University IACUC (protocol 10-003) and the temporary collection of salamanders was done under permit from the Ohio Department of Natural Resources.

LITERATURE CITED

- Bazar, M.A., M.J. Quinn Jr., K. Mozzachio, J.A. Bleiler, C.R. Archer, C.T. Phillips, and M.S. Johnson. 2010. Toxicological responses of Red-backed Salamander (*Plethodon cinereus*) exposed to aged and amended soils containing lead. Archives of Environmental Contamination and Toxicology 58:1040–1047.
- Bernal, M.H., K.R. Solomon, and G. Carrasquilla. 2009. Toxicity of formulated glyphosate (Glyphos) and Cosmo-Flux to larval and juvenile Colombian frogs 2. Field and laboratory microcosm acute toxicity. Journal of Toxicology and Environmental Health 72A:966– 973.
- Carlson, A.M., and D.L. Gorchov. 2004. Effects of herbicide on the invasive biennial *Alliaria petiolata* (garlic mustard) and initial responses of native plants in a southwestern Ohio forest. Restoration Ecology 12:559–567.
- Casper, G.S. 2005. *Plethodon cinereus* (Green, 1818).
 Pp. 796–800 *In* Amphibian Declines: The Conservation Status of United States Species. Lannoo, M. (Ed.) University of California Press, Berkeley, California, USA.
- Cole, E.C., W.C. McComb, M. Newton, C.L. Chambers, and J.P. Leeming. 1997. Response of amphibians to clearcutting, burning, and glyphosate application in the Oregon Coast Range. Journal of Wildlife Management 61:656–664.
- Dantzer, B.J., and R.G. Jaeger. 2007. Male Red-backed Salamanders can determine the reproductive status of conspecific females through volatile chemical signals. Herpetologica 63:176–183.
- Davic, R.D., and H.H. Welsh Jr. 2004. On the ecological roles of salamanders. Annual Review of Ecology, Evolution and Systematics 35:405–434.
- Dinehart, S.K., L.M. Smith, S.T. McMurry, T.A. Anderson, P.N. Smith, and D.A. Haukos. 2009. Toxicity of a glufosinate- and several glyphosatebased herbicides to juvenile amphibians from the Southern High Plains, USA. Science of the Total Environment 407:1065–1071.
- Gove, B., S.A. Power, G.P. Brinkley, and J. Ghazoul. 2007. Effects of herbicide spray drift and fertilizer overspread on selected species of woodland ground flora: comparison between short-term and long-term impact assessments and field surveys. Journal of Applied Ecology 44:374–384.
- Guynn, D.C., Jr, S.T. Guynn, T.B. Wigley, and D.A. Miller. 2004. Herbicides and forest biodiversity –

Wildlife Society Bulletin 32:1085-1092.

- Howe, C.M., M. Berrill, B.D. Pauli, C.C. Helbing, K. Werry, and N. Veldhoen. 2004. Toxicity of glyphosate-based pesticides to four North American frog species. Environmental Toxicology and Chemistry 23:1928-1938.
- Jones, B., S.F. Fox, D.M. Leslie, Jr., D.M. Engle, and R.L. Lochmiller. 2000. Herpetofaunal responses to brush management with herbicide and fire. Journal of Range Management 53:154-158.
- Maerz, J.C., N.L. Panebianco, and D.M. Madison. 2001. Effects of predator chemical cues and behavioral biorhythms on foraging activity of terrestrial salamanders. Journal of Chemical Ecology 27:1333-1344.
- Mann, R.M., and J.R. Bidwell. 2001. The acute toxicity of agricultural surfactants to the tadpoles of four Australian and two exotic frogs. Environmental Pollution 114:195-205.
- Mann, R.M., J.R. Bidwell, and M.J. Tyler. 2003. Toxicity of herbicide formulations to frogs and the implications for product registration: A case study from Western Australia. Applied Herpetology 1:13-22.
- Martin, S.B., R.G. Jaeger, and E.D. Prosen. 2005. Territorial Red-backed Salamanders can detect volatile pheromones from intruders. Herpetologica 61:29-35.
- McComb, B.C., L. Curtis, C.L. Chambers, M. Newton, and K. Bentson. 2008. Acute toxic hazard evaluations of glyphosate herbicide on terrestrial vertebrates of the Oregon Coast Range. Environmental Science and Pollution Research 15:266-272.
- Mushinsky, H.R., and E.D. Brodie Jr. 1975. Selection of substrate pH by salamanders. American Midland Naturalist 93:440-443.
- Relyea, R.A. 2005. The lethal impact of Roundup on aquatic and terrestrial amphibians. Ecological Applications 15:1118–1124.
- Semlitsch, R.D., C.A. Conner, D.J. Hocking, T.A.G. Rittenhouse, and E.B. Harper. 2008. Effects of timber harvesting on pond-breeding amphibian persistence: Testing the evacuation hypothesis. Ecological Applications 18:282–289.

- What do we know and where do we go from here? Shepard, J.P., J. Creighton, and H. Duzan. 2004. Forestry herbicides in the United States: an overview. Wildlife Society Bulletin 32:1020-1027.
 - Simmons, M.T., S. Windhager, P. Power, J. Lott, R.K. Lyons, and C. Schwope. 2007. Selective and nonselective control of invasive plants: The short-term effects of growing-season prescribed fire, herbicide, and mowing in two Texas prairies. Restoration Ecology 15:662-669.
 - Slaughter, B.S., W.W. Hochstedler, D.L. Gorchov, and A.M. Carlson. 2007. Response of Alliaria petiolata (Garlic Mustard) to five years of fall herbicide application in a southern Ohio deciduous forest. Journal of the Torrey Botanical Society 134:18-26.
 - Storrs Méndez, S.I., D.E. Tillitt, T.A.G. Rittenhouse, and R.D. Semlitsch. 2009. Behavioral response and kinetics of terrestrial atrazine exposure in American Toads (Bufo americanus). Archives of Environmental Contamination and Toxicology 57:590-597.
 - Sullivan, A.M., J.C. Maerz, and D.M. Madison. 2002. Anti-predator response of Red-backed Salamanders (Plethodon cinereus) to chemical cues from Garter Snakes (Thamnophis sirtalis): laboratory and field experiments. Behavioral Ecology and Sociobiology 51:227-233.
 - Takahashi, M. 2007. Oviposition site selection: Pesticide avoidance by Gray Treefrogs. Environmental Toxicology and Chemistry 26:1476-1480.
 - Vonesh, J.R., and J.C. Buck. 2007. Pesticide alters oviposition site selection in Gray Treefrogs. Oecologia 154:219-226.
 - Vonesh, J.R., and J.M. Kraus. 2009. Pesticide alters habitat selection and aquatic community composition. Oecologia 160:379-385.
 - Welsh, H.H., Jr., and S. Droege. 2001. A case for using plethodontid salamanders for monitoring biodiversity and ecosystem integrity of North American forests. Conservation Biology 15:558-569.
 - Zaffaroni, N.P., T. Zavanella, A. Cattaneo, and E. Arias. 1986. The toxicity of 2,4-dichlorophenoxyacetic acid to the adult Crested Newt. Environmental Research 41:79-87.



BENJAMIN GERTZOG (left) majored in biology at Denison University and is planning to attend graduate school in Physical Therapy. **LINDSAY KAPLAN** (center) also majored in biology at Denison and is looking to attend medical school in the future. **DAN NICHOLS** (right) was an economics major and biology minor at Denison who plans to attend graduate school in real estate development. (Unknown photographer)



GEOFF SMITH is an Associate Professor of biology at Denison University. He received his B.A. from Earlham College, and his Ph.D. from the University of Nebraska-Lincoln. His research examines the population and community ecology of reptiles and amphibians, and in particular, how human-induced changes in the environment are affecting them. (Photographed by James Hale)



JESSICA RETTIG is an Associate Professor of biology at Denison Unversity. She received her B.A. from Earlham College, and her Ph.D. from Michigan State University. Her research focuses on the community ecology of small temperate ponds, and in particular the ecology of Bluegill Sunfish (*Lepomis macrochirus*). (Photographed by James Hale)