# **R**EPTILES TRADED IN THE FETISH MARKET OF LOMÉ, TOGO (WEST AFRICA)

GABRIEL H. SEGNIAGBETO<sup>1,2</sup>, FABIO PETROZZI<sup>3</sup>, AURÉLIE AÏDAM<sup>4</sup>, AND LUCA LUISELLI<sup>3,5</sup>

<sup>1</sup>Département de Zoologie et Biologie Animale, Faculté des Sciences, Université de Lomé, BP. 1515, Lomé, Togo <sup>2</sup>AGBO-ZEGUE NGO (Togolese Society for Nature Conservation), BP. 6057 Lomé, Togo, e-mail: h\_segniagbeto@yahoo.fr <sup>3</sup>Centre of Environmental Studies Demetra, Rome, Italy, e-mail: fapetrozzi@gmail.com

<sup>4</sup>Centre d'Etudes Biologiques de Chizé, CNRS, UPR 1934, Villiers en Bois 79360, Beauvoir sur Niort, France, e-mail:

aurelie.aidam@gmail.com

<sup>5</sup>Corresponding author, e-mail: lucamlu@tin.it

*Abstract.*—Fetish beliefs are important to the culture of West African people. Many animals, including reptiles, are traded in fetish markets and used for traditional medicine and for their perceived magical properties. There are few available quantitative surveys of the reptiles traded in such fetish markets in the international literature. Here, we report the results of a survey of the reptiles traded in the largest fetish market of West Africa (situated in Lomé, Togo). The market was visited in three distinct periods (July 2012, December 2012, and February 2013), but counts of all traded specimens were done only in December 2012. A total of 1,765 reptile individuals, belonging to 37 different species, were recorded. The great majority of species came from two vegetation zones (forest and Guinea savannah), with the great majority of individuals coming from the Guinea savannah region. Chameleons (*Chamaeleo senegalensis*) dominated in the sample. Among snakes, the Puff Adder (*Bitis arietans*) was frequently traded. The conservation implications of the observed pattern are discussed.

Key Words.-conservation; fetish market; Reptilia; Togo; West Africa

#### INTRODUCTION

Trade in animals for purposes other than for food or as pets is on the increase. The Chinese medicine trade and its impacts on wildlife are particularly well known (e.g., Bauer 2009), including its impacts on African wildlife (rhino horns especially; Leader-Williams 1992). Reptiles are often of particular focus (Simelane and Kerley 1997, 1998); for example, Bauer 2009 addresses the depletion of Tokay Geckos (Gekko gecko) in southeastern Asia for the production of potions. In Asia, the economic boom has obviously seen an increase in the demand for what were previously small-market trinkets, traditional medicines, and what one may call a "fetish". Indeed, a 'fetish' is defined as an animal, plant, or mineral-derived part that is used in traditional religion or traditional medicine practices (Mockler-Ferryman 1898; Nikolaus 2011). Thus, fetish markets are different from bushmeat markets in scope and type of traded animals, given that the latter ones serve the scope of domestic consumption for food (e.g., Fa et al. 2003; van Vliet et al. 2012); whereas, the former ones are more tightly linked to perceived magic and traditional cultural practices (Taylor and Fox 1992).

Traditional medicine and fetish beliefs are important for the culture of African people, especially in the central (Gilli 1987; Pauwels et al. 2003), southern (de Surgy 1993; Whiting et al. 2011), and western (Adeola 1992; Fretey et al. 2007) countries where there is a flourishing market of animals and animal parts. Among the many animal taxa observed in these markets (e.g., birds, mammals, amphibians, invertebrates), reptiles are typically present (Simelane and Kerley 1997, 1998; Fa et al. 2003; van Vliet et al. 2012).

In spite of the scattered information found in the available literature on the reptiles species traded, few quantitative studies have been performed on reptiles traded in single fetish markets (but see Fretey et al. 2007). Collection of this type of quantitative data is important in terms of conservation biology because several species of West African reptiles are declining considerably (Luiselli et al. 2007; Reading et al. 2010). For instance, it is important to understand whether particular communities make use of certain groups of organisms, and in which habitats or seasons they catch these animals (Adeola and Decker 1987; Edwards 2003; Willcox and Nambu 2006).

In this paper, we quantify the reptile species

traded in a single fetish market of West Africa. We studied the "Marché au Fétiches" (French for fetish market) situated in Lomé, the capital city of Togo. This market is particularly interesting, compared to other similar marketplaces in West African countries, because it is considerably larger in size (thus including a higher number of specimens traded) and because it is situated in a town surrounded by areas with different vegetation types (Guinea savannah, forest, coastal sites, etc.; Ern 1979) that may influence the types of reptiles captured.

## **MATERIALS AND METHODS**

The fetish market is situated in Akodessewa. in the east of Lomé, Togo. The market was moved to the present site at the end of the 1990s, with the purpose of creating a dedicated venue for selling traditional medicine away from the main Bè market. This is the most famous fetish market of West Africa because it offers a large choice to customers who come looking for the ingredients of sorcerer's and traditional healers' prescriptions.

We surveyed the market place three times. First, one of the authors (LL) conducted a preliminary survey in July 2012 to determine the feasibility of this study, with identification of the species traded and with no count of the number of traded specimens (apart from chelonians and pythons whose corpses were counted). Second, we conducted a quantitative survey in December 2012. We examined and counted all of the individual reptile specimens available at the marketplace. The number of individuals traded by species was determined by recording the carcasses, heads, and body parts clearly attributable to different specimens. Third, we conducted another survey by recording of all species traded in February 2013. Counts were simultaneously conducted by four authors, each of us analyzing a distinct portion of the market stands, and overall we spent a total of 34 personhours of field effort. We could not exclude the possibility that minor errors in determining the exact number of individuals of all species may have occurred. However, we used the most conservative criteria for determining the numbers of individuals, although not all species market was also assigned to a pertinent

would be affected in the same way by this approach. For instance, if for a given snake species we found ten different heads and ten different skins of consistent size, then we considered the number of traded specimens to be 10. This conservative approach might have underestimated the number of snakes, whereas for lizards, which were present in the market as whole specimens, the number was likely to be close to the full number of specimens present.

In each of the three surveys, we walked up to the market trader and openly identified corpses and counted skins, heads, and other body parts. We also spoke to the traders in order to obtain more information on the animals traded. including their site of capture if known. In our study market, traders were disinclined to allow people to make notes on their products for fear of interference by government officials, as may happen in other similar marketplaces (Luiselli et al., pers. obs.). For this reason, by explaining our work, we made sure that traders did not hide their products away during surveys.

Each specimen was identified to species level (using Chippaux 2006 for snakes, and Trape et al. 2012 for lizards, turtles, tortoises, and crocodiles), although this was not possible for some cases (e.g., Meizodon spp.). In addition, given the confused taxonomic status of the genus *Psammophis* (cf. Hughes 1999), we assigned those specimens fitting with 'phillipsii' or 'sibilans' according to the identification keys of Chippaux (2006). Some species were recently subjected to splitting, with West African populations being considered distinct taxa from other African populations. This was the case, for instance, of Bitis gabonica to Bitis rhinoceros (Lenk et al. 1999), Kinixys belliana to Kinixys noguevi (Lataste 1886; Rhodin et al. 2012), and Crocodylus niloticus to Crocodylus suchus (Schmitz et al. 2003). Here we maintain both nomenclatures, placing the 'old' taxon in parentheses (e.g., *Kinixys* [*belliana*] *nogueyi*). Where assessed, the International Union for Conservation of Nature (IUCN) status of the each species was noted (IUCN Red List of Threatened Species. Available from http://www.iucnredlist.org [Accessed 21 December 2012]). Each species observed in the

## Herpetological Conservation and Biology

vegetation zone based on its distribution range region according to White (1963). Thus, the across West Africa. Maps used for this assignment were taken from IUCN 2012 (op. *cit.*) when available, or from Chippaux (2006) and Trape et al. (2012) in all other cases. The territory of Togo was subdivided into five main vegetation zones (Ern 1979): (1) northern plains (= transition area between Sudanese-Sahelian savannahs); (2) hilly dry woodlands and savannah mosaics (Sudanese savannah); (3) Guinean savannah; (4) Togo mountains forests (= semi-deciduous moist forest); and (5)southern coastal plain. In our case, we simplified this categorization by condensing vegetation zones (1) and (2) into the novel category 'dry savannah' because both of these zones were basically belonging to the Sudanese savannah

following vegetation zone categories were considered for our analyses: (1) dry savannah, (2) Guinean savannah; (3) forest; (4) marine/coastal; and (5) generalist (when more than two vegetation zones were inhabited per given species).

### RESULTS

In total, 1,765 individual reptiles, belonging to 37 different species, were observed (Table 1; Fig. 1). Sea turtles were traded as heads or shells, tortoise and freshwater turtles were dried entire specimens or shells, and snakes were either heads, skins, or dried specimens. Snake bones were also observed, but in these cases the species remained unidentified. Lizards were dried entire



FIGURE 1. Examples of the reptile species displayed for sale at the fetish market of Lomé, Togo. Chameleons, and especially *Chamaeleo senegalensis*, dominated among the reptiles species traded (a). Among snakes, the Puff Adder (Bitis arietans) was the most heavily traded species (b). In this photo, together with many Puff Adder heads, there are also some heads of the Forest Cobra (Naja melanoleuca). Tortoises were not heavily traded, but Kinixys (belliana) nogueyi was the most commonly traded species (c). Skins of crocodiles were also observed in the market (d). Photo d shows Crocodylus (niloticus) suchus. All photos by Aurélie Aïdam.

**TABLE 1.** Number and species identity of reptile specimens censured at the Fetish market, Lomé in December 2012, including their main habitat/vegetation zone of origin. Nomenclature follows Chippaux (2006) for snakes, and Trape et al. (2012) for lizards, turtles, tortoises, and crocodiles.

Species name	Common name	No. in market	Forest	Guinea savannah	Dry savannah	Marine/ coastal
Crocodilians						
Crocodylus (niloticus) suchus	West African Nile Crocodile	54	1	1	1	0
Mecistops (= Crocodylus) cataphractus	African Slender-snouted Crocodile	1	1	0	0	0
Testudines						
Pelomedusa subrufa	African Helmeted Turtle	2	0	1	0	0
Pelusios castaneus	Western Hinged Terrapin	27	0	1	0	0
Centrochelys (= Geochelone) sulcata	African Spurred Tortoise	2	0	0	1	0
Kinixys (belliana) nogueyi	Bell's Hinged Tortoise	16	0	1	0	0
Kinixys homeana	Home's Hinged Tortoise	1	1	0	0	0
Kinixys erosa	Forest Hinged Tortoise	4	1	0	0	0
Lepidochelys olivacea	Olive Ridley Sea Turtle	6	0	0	0	1
Chelonia mydas	Green Sea Turtle	1	0	0	0	1
Dermochelys coriacea	Leatherback sea turtle	1	0	0	0	1
Trionyx triunguis	Nile Soft-shelled Terrapin	2	1	0	0	1
Cyclanorbis senegalensis	Senegal Flapshell Terrapin	1	0	1	0	0
Sauria						
Agama agama	Common Rainbow Lizard	128	1	1	1	0
Chamaeleo gracilis	Graceful Chameleon	29	1	1	0	0
Chamaeleo senegalensis	Senegal Chameleon	963	0	1	1	0
Trachylepis perrotetii	Spotted Savannah Skink	6	0	1	0	0
Varanus ornatus	Ornate Monitor	1	1	0	0	0
Varanus niloticus	Nile Monitor	31	0	1	0	0
Varanus exanthematicus	Western Savannah Monitor	9	0	0	1	0
Serpentes						
Python sebae	African Rock Python	28	1	1	0	0
Python regius	Ball Python	16	1	1	0	0
Toxicodryas blandingii	Powdered Tree Snake	6	1	0	0	0
Psammophis cf. phillipsii	Olive Whip Snake	25	1	0	0	0
Psammophis cf. sibilans	Short-snouted Sand Snake	12	0	1	1	0
Thelotornis kirtlandii	Forest Vine Snake	1	1	0	0	0
Philothamnus irregularis	Irregular Green Snake	11	1	1	0	0
Hapsidophrys smaragdina	Emerald Snake	1	1	0	0	0
Lamprophis fuliginosus	Brown House Snake	3	1	1	0	1
Meizodon sp.	Crowned Snake	1	0	1	1	0
Rhamphiophis oxyrhynchus	Western Beaked Snake	21	0	1	1	0
Dasypeltis gansi	Gans' Egg-eating Snake	1	0	1	1	0
Naja melanoleuca	Forest Cobra	34	1	1	0	0
Naja nigricollis	Spitting Cobra	5	1	1	0	1
Dendroaspis viridis	West African Green Mamba	4	1	0	0	0
Bitis arietans	Puff Adder	257	0	1	1	1
Bitis (gabonica) rhinoceros	Gaboon Viper	21	1	0	0	0
Undetermined	-	33				
TOTAL		1,765	16	16	8	4

specimens.

There was an uneven distribution of number of species in terms of their main habitat of origin, with the great majority of species coming from two vegetation zones (forest and Guinea savannah; Table 1). Species from dry savannah, generalist, and marine/coastal species were under-represented in the sample. The majority of specimens traded were Guinean savannah species (79.2%), compared with forest species (11.4%; Table 1). In this case, savannah specimens were clearly more traded than specimens from any other vegetation zone.

Chameleons (*Chamaeleo senegalensis*) were by far the most intensively traded reptiles, accounting for nearly 55% of the total number of reptiles examined (Table 1). These chameleons are typically savannah inhabitants (Chirio and LeBreton 2007; Trape et al. 2012). The second most commonly traded species was the Puff Adder (*Bitis arietans*), also a savannah species (Chippaux 2006; Chirio and LeBreton 2007).

In terms of IUCN conservation status, one species was Critically Endangered (the Leatherback Sea Turtle, *Dermochelys coriacea*), one was Endangered (the Green Sea Turtle, *Chelonia mydas*), three were Vulnerable (African Spurred Tortoise, *Centrochelys* (= *Geochelone*) *sulcata*; Home's Hingeback Tortoise,*Kinixys homeana*; and Olive Ridley Sea Turtle, *Lepidochelys olivacea*), and one was Lower Risk/Near Threatened (Senegal Flapshell Turtle, *Cyclanorbis senegalensis*). All other species were either not listed, Least Concern, or Data Deficient.

## DISCUSSION

Overall, our study revealed that a high number of reptile individuals, from a varied diversity of species, are currently traded in the Lomé fetish market. The number of reptiles species identified in this study (n = 37) is not comparable to that identified in a previous survey of the same marketplace (Taylor and Fox 1992) because Taylor and Fox reported only the presence of such general groups as crocodiles, snakes, sea turtles, turtles, and chameleons. The reptiles observed in the present surveys were mainly

from the Guinean savannah habitat, which is the predominant vegetation type just around Lomé and in most part of Togo (Ern 1979). Therefore, we suggest that the majority of the specimens used for the traditional medicine and fetish were collected opportunistically close to the site where they were actually marketed. For instance, in the February 2013 survey, we observed 22 freshly killed (not yet dried) snake corpses (7 Naja nigricollis, 12 Bitis arietans, 3 Psammophis cf. *phillipsii*) that, according to their sellers, were captured along the eastern peripheral side of Lomé. However, there were certainly exceptions to this rule. For instance, a specimen of the tortoise *Centrochelys sulcata* was present in the market despite being absent from Togo (Trape et al. 2012), and we also observed in the Lomé market the skin of a Lion (Panthera leo *senegalensis*) which was reportedly originating from Pendjari Biodiversity Reserve (Benin). However, both giant tortoises and lions are charismatic animals for the local cultures (Schaller 1972; Chardonnet 2002; Branch 2007) and it is not surprising that the demand can be so high for such species that collectors may make long displacements to bring them to the marketplace. All the other species of traded reptiles are known to occur in Togo and in the surrounding area of Ghana (Hughes and Barry 1969; Leaché 2005) and Benin (Leaché et al. 2006; Segniagbeto 2009; Ullenbruch et al. 2010; Segniagbeto et al. 2011). Concerning *Centrochelys sulcata*, although no natural populations are currently known in Togo (Branch 2007), this species is ranched in southern Togo for the international pet trade export (Ineich 2011), Togo being the main export country for Africa (Convention of International Trade on Endangered Species. 1980. CITES database analysis. Available from http://www.cites.org/eng/resources/species.html [Accessed September 2012].

It is noteworthy that, at least for some species, there is a tight link between the bushmeat market and the fetish market (Fa et al. 2003). For instance, some of the larger animals are mainly sold in parts (crocodiles, pythons, large chelonians), with the meat being used for the food trade and the bones, heads or skins being used for fetish purposes. This potential link

between bushmeat and fetish markets should be explored more in depth by further studies. Certainly, the use of reptile body parts (fat, skull, bone, organs, and blood) is very specific for fetish practices, as each body part is considered to be essential in traditional pharmacopoeia for various diseases (e.g., convulsions, malaria, fever, HIV-AIDS, liver problems, tetanus, induce vomiting, etc.), although of course no scientific evidence of any benefit from these animal parts has ever been demonstrated in the medicinal literature. Specific body parts are also used for additional reasons other than healthcare. For instance, many businessmen in Togo believe that appropriate animist ceremonies using reptile body parts can help them to improve their activities, and to this purpose they come regularly to ask the advice of a voodoo priest in this market. It is also common to see reptilian body parts in the small statuettes of voodoo in the fetish markets.

Most of the reptile species observed in the market are of lesser international conservation concern, while those of high conservation concern were observed in very low numbers (e.g., sea turtles: Dermochelys coriacea and *Chelonia mydas*). It is possible that the scarcity of threatened species in the market may reflect their disappearance in the field, as already demonstrated to occur in bushmeat markets where the species threatened by hunting always become rare in markets after the maximum sustainable yield has been overshot (Cowlishow et al. 2005). In addition, some of the traded species are declining locally or even regionally (e.g., Kinixys homeana and Kinixys erosa, Luiselli 2003a, 2003b; Luiselli et al. 2006; Python sebae, Luiselli et al., 2007; Python regius, Reading et al. 2010; and Mecistops (Crocodylus) cataphractus, Chirio and LeBreton 2007; Shirley 2010), and the overall IUCN Red List assessment for many of these species is currently in preparation. The trade of these declining species warrants monitoring by competent agencies, with special attention on whether any type of cross-border trade (e.g., from Ghana and Benin) might occur to refurnish the Lomé fetish market. Indeed, during the '70s and '80s, Lomé was well-known for selling (and even exporting to Europe) many reptile these markets for the interests of people and

specimens which were originally captured in Ghana (Laurent Chirio and Barry Hughes, pers. comm. to LL), and it is possible that such an international trade may still occur on some extent (Ineich 2011).

Despite being utilized for fetish purposes, several of the reptile species recorded in the Lomé market are legally protected in Togo under Article 62, Section 2, Chapter II of the law N° 2008-005, regulating environment protection and wildlife conservation. This is the case for the four marine turtle species (Dermochelys coriacea, Lepidochelys olivacea, Chelonia mydas, and Eretmochelys imbricata), for all of the freshwater turtles and tortoises, for crocodiles, and for Python sebae. In the practice, however, it is very difficult for the administration in charge of wildlife conservation to implement this law, because most Togolese people think that all reptiles, and in particular snakes, are dangerous (and thus that they should be killed), and/or are useful as meat and for traditional medicine. Indeed, marine turtles, freshwater turtles, tortoises, African Rock Pythons and monitor lizards are considered as game and are systematically hunted (Segniagbeto et al., pers. obs.).

A shortcoming of this study is that we provided only a snapshot of the current reptiles traded as fetish in Togo because we surveyed the market for only a short time span. For instance, the number and even the species of traded chelonians varied between the first and the second survey (see the data concerning the number of chelonians recorded in the first survey in Luiselli et al. (In press). Therefore, longitudinal monitoring of the reptiles traded in the Lomé fetish market may allow much more detailed understanding of the trade dynamics of the targeted species. In addition, research on fetish markets, as with research on bushmeat markets, should aim to understand: (1) the history of the trade; (2) the perspectives of those involved; (3) the socio-economic background of the marketplace and community; (4) the biotic background of the specimens; and (5) the spatial and temporal patterns of species being traded. This information is crucial to make appropriate assessments and decisions on how to manage

wildlife alike.

Acknowledgments.—We thank Dr. Jamie Carr (UNEP, Cambridge) and Prof. John F. Oates (CUNY, New York) for having participated in the first survey of the market together with one of the authors (LL), and Dr. Daniele Dendi for having helped us in measuring the traded specimens. We also thank Dr. Barry Hughes (London), Dr. Laurent Chirio (Libreville, Gabon), Mr. Gbodjivi Pascal (Lomé), Mr. Limikpo Jacob (President of the Fetish Market, Lomé), and Dr. Jean-Francois Trape (IRD, Dakar) for useful information. This research was supported by funds from the Associazione Centro Elis (Rome, Italy; funds to LL and FP).

## LITERATURE CITED

- Adeola, M.O. 1992. Importance of wild animals and their parts in the culture's religious festivals, and traditional medicine, of Nigeria. Environmental Conservation 19:125–134.
- Adeola, M.O., and E. Decker. 1987. Wildlife utilization in rural Nigeria. Pp. 512–521 *In* Proceedings of the International Symposium and Conference on Wildlife Management in Sub-Saharan Africa. Clers, B.D. (Ed.). Harare, Zimbabwe.
- Bauer, A.M. 2009. Geckos in traditional medicine: forensic implications. Applied Herpetology 6:81–96.
- Branch, B. 2007. Tortoises, Terrapins, and Turtles of Africa. Struik Publishers, Cape Town, South Africa.
- Chardonnet, P. (Ed.). 2002. Conservation of the African Lion: Contribution to a Status Survey. International Foundation for the Conservation of Wildlife, France and Conservation Force, USA.
- Chippaux, J.P. 2006. Les serpents d'Afrique occidentale et centrale. Collection Faune et Flore Tropicales, IRD Editions, Paris, France.
- Chirio, L., and M. LeBreton. 2007. Atlas des reptiles du Cameroun. Collection Patrimoines naturels n°67, Muséum national d'Histoire naturelle, IRD Editions, Paris, France.
- Cowlishaw, G., S. Mendelson, and J. Rowcliffe. 2005. Evidence for post depletion

sustainability in a mature bushmeat market. Journal of Applied Ecology 42: 460–468

- *Acknowledgments.*—We thank Dr. Jamie Carr JNEP, Cambridge) and Prof. John F. Oates CUNY, New York) for having participated in
  - Edwards, I.B. 2003. The fetish market and animal parts trade of Mali, West Africa: an ethnographic investigation into cultural use and significance. Ph.D. Dissertation, Oregon State University, Corvallis, Oregon, USA. 231 p.
  - Ern, H. 1979. Vegetation Togos. Gliederung, Gefährdung, erhaltung. willdenowia 9:295– 312.
  - Fa, J.E., D. Curries, and J. Meeuwig. 2003. Bushmeat and food security in the Congo Basin: linkages between wildlife and people's future. Environmental Conservation 30:71–78.
  - Fretey, J., G.H. Segniagbeto, and M. Souma. 2007. Presence of sea turtles in traditional pharmacopoeia and beliefs of West Africa. Marine Turtle Newsletter 116:23–25.
  - Gilli, B. 1987. Heviesso et le bon ordre du monde. Lomé, éd. Haho.
  - Hughes, B. 1999. Critical review of a revision of *Psammophis* (Linnaeus, 1758) (Serpentes, Reptilia) by Frank Brandstaetter. African Journal of Herpetology 48:63–70.
  - Hughes, B., and D.H. Barry. 1969. The snakes of Ghana: a checklist and key. Bulletin de l'Institut fundamental d'Afrique Noire 31:1004–1041.
  - Ineich, I. 2011. Les élevages de reptiles et de scorpions au Benin, Togo et Ghana – plus particulièrement a gestion des quotas d'exportation et la définition des codes "source" des spécimens exportes. Projet CITES A-251, Berne, Switzerland.
  - Lataste, F. 1886. Description d'une tortue nouvelle du Haut-Sénégal (*Homopus nogueyi*). Le Naturaliste 2:286–287.
  - Leaché, A. D. 2005. Results of a herpetological survey in Ghana and a new country record. Herpetological Review 36:16–19.
  - Leaché, A.D., M.O. Rödel, C.W. Linkem, R.E. Diaz, A. Hillers, and M.K. Fuijita. 2006. Biodiversity in a forest island: reptiles and amphibians of the West African Togo hills. Amphibian and Reptile Conservation 4:22–45.
  - Leader-Williams, N. 1992. The world trade in

rhino horn: a review. Traffic International, World Wildlife Federation, Cambridge, UK.

- Lenk, P., H.W. Hermann, and U. Joger. 1999. Phylogeny and taxonomic subdivision of *Bitis* (Reptilia, Viperidae) based on molecular evidence. Kaupia 8:31–38.
- Luiselli, L. 2003a. Assessing the impact of human hunting activities on populations of forest tortoises (genus *Kinixys*) in the Niger Delta, Nigeria. Chelonian Conservation and Biology 4:735–738.
- Luiselli, L. 2003b. Comparative abundance and population structure of sympatric Afrotropical tortoises in six rainforest areas: the differential effects of "traditional veneration" and of "subsistence hunting" by local people. Acta Oecologica 24:157–163.
- Luiselli, L., F. Petrozzi, and G.C. Akani. In press. Long-term comparison reveals trends in turtle trade in bushmeat markets of southern Nigeria. Herpetozoa.
- Luiselli, L., E. Politano, and J. Lea. 2006. Assessment of Vulnerable status of *Kinixys homeana* (Testudines: Testudinidae) for the IUCN Red List. Chelonian Conservation and Biology 5:130–139.
- Luiselli, L., G.C. Akani, E.A. Eniang, and E. Politano. 2007. Comparative ecology and ecological modeling of sympatric pythons, *Python regius* and *P. sebae*. Pp 88–100. *In* Biology of the Boas and Pythons. Henderson, R.W., and R. Powell, (Eds). CPG/Biological Science Press, New York, New York, USA.
- Mockler-Ferryman, A.F. 1898. Imperial Africa: The Rise, Progress and Future of the British Possessions in Africa. Volume 1. Imperial Press, London, UK.
- Nikolaus, G. 2011. The fetish culture in West Africa: an ancient tradition as a threat to endangered birdlife? Bonner Zoologische Monographien 57:145–149.
- Pauwels, O.S.G., M.O. Rödel, and A.K. Toham. 2003. *Leptopelisnotatus* (Anura: Hyperoliidae) in the Massif du Chaillu, Gabon: from ethnic wars to soccer. Hamadryad 27:271–273.
- Reading, C. J., L.M. Luiselli, G.C. Akani, X. Bonnet, G. Amori, J.M. Ballouard, E. Filippi, G. Naulleau, D. Pearson, and L. Rugiero. 2010. Are snake populations in widespread decline? Biology Letters 6:777–780.

- Rhodin, A.G.J., P.P. Van Dijk, J.B. Iverson, and H.B. Shaffer. 2012. Turtles of the world, 2012 update: annotated checklist of taxonomy, synonymy, distribution, and conservation status. Turtle Taxonomy Working Group. Pp. 000.85–000.164 In Conservation Biology of Turtles and Tortoises: Freshwater а Compilation Project of the IUCN/SSC Tortoise and Freshwater Turtles Specialist Group. Rhodin, A.G.J., P.C.H. Pritchard, P.P. Van Dijk, R.A. Saumure, K.A. Buhlmann, J.B. Iverson, and R.A. Mittermeier (Eds.). Chelonian Research Monographs, Lunenburg, USA.
- Schaller, G.B. 1972. The Serengeti Lion. University of Chicago Press, Chicago, Illinois, USA.
- Schmitz, A., P. Mansfeld, E. Hekkala, T. Shine,
  H. Nickel, G. Amato, and W. Boehme. 2003.
  Molecular evidence for species level divergence in African Nile Crocodiles *Crocodylus niloticus* (Laurenti, 1768).
  Comptes Rendus Palevolution 2:703–712.
- Segniagbeto, G.H. 2009. Herpétofaune du Togo: taxinomie, biogéographie. Ph.D. Dissertation, University of Lomé (Togo) and National Museum of Natural History, Paris, France. Vol. I: 1–172 & Vol. II: 1–192.
- Segniagbeto, G.H., J.F. Trape, P. David, A. Ohler, A. Dubois, I.A. Glitho. 2011. The snake fauna of Togo: systematics, distribution and biogeography, with remarks on selected taxonomic problems. Zoosystema 33:325–360.
- Shirley, M.H. 2010. Slender-snouted Crocodile *Crocodylus cataphractus*. Pp. 54–58 *In* Crocodiles Status Survey and Conservation Action Plan. 3<sup>rd</sup> Edition. Manolis, S.C., and C. Stevenson (Eds.). Crocodile Specialist Group, Darwin, Australia.
- Simelane, T.S., and G.I.H. Kerley. 1997. Recognition of reptiles by Xhosa and Zulu communities in South Africa, with notes on traditional beliefs and uses. African Journal of Herpetology 46:49–53.
- Simelane, T.S., and G.I.H. Kerley. 1998. Conservation implications of the use of vertebrates by Xhosa traditional healers in South Africa. South African Journal of Wildlife Research 28:121–126.
- Taylor, M.C., and J. Fox. 1992. The fetish market, Lomé, Togo. The Nigerian Field

## 57:119–125.

- Trape, J.F., L. Chirio, and S. Trape. 2012. Lézards, crocodiles et tortues d'Afrique occidentale et du Sahara. IRD Editions, Paris.
- Ullenbruch, K., O. Grell, and W. Böhme. 2010. Reptiles from southern Benin, West Africa, with the description of a new *Hemidactylus* (Gekkonidae), and a country-wide checklist. Bonn Zoological Bulletin 57:31–54.
- van Vliet, N., C. Nebesse, S. Gabalemoke, D. Akaibe, and R. Nasi. 2012. The bushmeat market in Kisangai, Democratic Republic of Congo: implications for conservation and food









security. Oryx 46:196–203.

- White, F. 1963. The Vegetation of Africa. UNESCO Press, Paris, France.
- Whiting, M.J., V.L. Williams, and T.J. Hibbitts. 2011. Animals traded for traditional medicine at the Faraday market in South Africa: species diversity and conservation implications. Journal of Zoology 284:84–96.
- Willcox, A.S., and D.M. Nambu. 2006. Wildlife hunting practice and bushmeat dynamics of the Banyangi and Mbo people of south-western Cameroon. Biological Conservation 134:251– 261.

**GABRIEL H. SEGNIAGBETO**, of Togolese nationality, currently teaches Zoology at the University of Lomé (Togo) and leads the AGBO-ZEGUE ONG (Togolese Society for Nature Conservation), also based in Lomé. He is an expert of the taxonomy of Togolese reptiles and amphibians, and is currently doing research on the ecology and conservation of West African reptiles. He has been co-author of the two most recent and comprehensive studies on the snakes and amphibians of Togo, and is currently working on a similar review focused on Togolese lizards. (Photographed by Fabrio Petrozzi).

**FABIO PETROZZI**, of Italian nationality, is an ecologist and wildlife management expert who has been working on conservation and ecology projects in West Africa (especially in Ghana and Togo) for several years on behalf of the Ministry of Foreign Affairs of the Republic of Italy. Apart from West African ecological herpetology, his main scientific interests are in sustainable development and applied ecology in savannah and forest ecosystems of West Africa, with special focus on the bushmeat trade analysis and on the conservation of the Lion (*Panthera leo*). (Photographed by Massimiliano Di Vittorio).

AURÉLIE AÏDAM, of French and Togolese nationalities, has been working at the CNRS, France and is now also working with Togo Faune. She is an expert of snake ecophysiology. In particular, she has studied the effects of temperature on the early postnatal growth of *Vipera aspis* and the demography, natural history, and color polymorphism of the European Dice Snake, *Natrix tessellata*. In Togo, she is also investigating the ecology, ecotoxicology, and conservation biology of native snakes, mainly pythons. (Photographed by Gabriel Segniabeto).

LUCA LUISELLI, of Italian nationality, is a member of the Steering Committee of IUCN/SSC TFTSG and the Associate Editor of several international journals such as Amphibia-Reptilia, Endangered Species Research, the Open Ecology Journal, and Chelonian Conservation and Biology. He is author of over 150 international scientific papers, several of them published in top journals (Nature, Ecology, Oikos, Oecologia, Biology Letters, Chemosphere, Biological Conservation, etc.). He has been working in West Africa (particularly Nigeria) for 17 years as a project director for several ecological projects led by such organizations as ENI and IUCN/SSC TFTSG, and also teaches Ecological Statistics at the University 'Roma Tre'. He has won 11 international research grants (e.g., the Mohamed Bin Zayed Conservation Fund). His main research interests are in tropical forest ecology, community ecology, theoretical ecology, and conservation biology of reptiles, especially snakes and chelonians. (Photographed by Fabio Petrozzi).