

---

## INTRA- AND INTER-POPULATION VARIABILITY OF FOOD PREFERENCES OF TWO *Natrix* SPECIES ON THE BALKAN PENINSULA

GORAN ŠUKALO<sup>1</sup>, SONJA ĐORĐEVIĆ<sup>2,5</sup>, SLADANA GVOZDENOVIC<sup>3</sup>, ALEKSANDAR SIMOVIĆ<sup>2,5</sup>, MARKO ANĐELKOVIĆ<sup>4,5</sup>, VELJKO BLAGOJEVIĆ<sup>4</sup>, AND LJILJANA TOMOVIĆ<sup>2,4,5</sup>

<sup>1</sup>University of Banja Luka, Faculty of Sciences, Mladena Stojanovića 2, 78000 Banja Luka, Republic of Srpska, Bosnia and Herzegovina; e-mail: sukalogoran@yahoo.com

<sup>2</sup>University of Belgrade, Faculty of Biology, Institute of Zoology, Studentski trg 16, 11000 Belgrade, Serbia, e-mails: sonjadj@bio.bg.ac.rs, lili@bio.bg.ac.rs

<sup>3</sup>University of Montenegro, Institute for Marine Biology, Dobrota b.b., P.O. Box 69 85330 Kotor, Montenegro, e-mail: sladjana87gvozdencovic@yahoo.com

<sup>4</sup>University of Belgrade, Institute for Biological Research “Siniša Stanković,” Department of Evolutionary Biology, Bulevar despota Stefana 142, 11000 Belgrade, Serbia, e-mails: marko.andjelkovic@ibiss.bg.ac.rs, veljko.blagojevic1988@gmail.com

<sup>5</sup>Serbian Herpetological Society “Milutin Radovanović,” Bulevar despota Stefana 142, 11000 Belgrade, Serbia, e-mail: alexandar.simovic@gmail.com

**Abstract.**—In aquatic and surrounding terrestrial ecosystems, snakes of the genus *Natrix* are among the top predators, feeding predominantly on fishes and amphibians, but also on other reptiles and small mammals. In the diets of *Natrix natrix* and *N. tessellata*, preferred food items vary geographically and during ontogeny. To understand these variations, we collected data on their diet composition in several habitats (river, wetlands, and two types of lakes), some of which are under severe anthropogenic pressures. Both *Natrix* species were able to quickly adapt to changes in available prey, and to feed even on non-indigenous, invasive, and potentially hazardous fish. Disturbed ecosystems are particularly susceptible to invasions. Therefore, if alien fishes become dominant, they may threaten populations of native snakes. On the other hand, semi-aquatic snakes can contribute to natural regulation of alien fish species abundance. The preliminary results presented herein emphasize the urgent need for integrative studies of natricine snakes and their prey ecology under various and/or dynamic circumstances. Better understanding of the functioning of different aquatic ecosystems may enable proper conservation and restoration for these snake species and their habitats.

**Sažetak.**—U akvatičnim i okolnim terestričnim ekosistemima, zmije iz roda *Natrix* često su pri vrhu lanaca ishrane. Hrane se pretežno ribom i vodozemcima, ali i drugim gmizavcima i sitnim sisarima. Omiljeni plijen *Natrix natrix* i *N. tessellata* varira, kako geografski tako i tokom ontogenije. Kako bismo mogli razumjeti navedene razlike, prikupili smo podatke o sastavu ishrane ovih zmija u nekoliko staništa (rijeka, močvarna staništa i dva tipa jezera); neka od navedenih područja pod intenzivnim su antropogenim pritiscima. Objе vrste roda *Natrix* sposobne su se brzo prilagoditi promjenama u dostupnom plijenu, i hraniti se čak i alohtonim, invazivnim i potencijalno opasnim vrstama riba. Narušeni ekosistemi naročito su podložni naseljavanju invazivnim vrstama. U tom smislu, ukoliko bi strane vrste riba postale dominantne, mogle bi ugroziti populacije autohtonih vrsta zmija. Sa druge strane, semiakvatične vrste zmija mogu doprinijeti prirodnoj regulaciji brojnosti alohtonih vrsta riba. Preliminarni rezultati koje ovdje prikazujemo naglašavaju potrebu da se u najskorije vrijeme sprovedu integrativne studije ekologije akvatičnih zmija i njihovog plijena pod različitim i/ili promijenljivim okolnostima. Bolje poznavanje funkcionisanja različitih akvatičnih ekosistema može omogućiti adekvatnu zaštitu i oporavak navedenih vrsta zmija i njihovih staništa.

**Key Words.**—Dice Snakes; diet shifts; disturbed habitats; Grass Snakes; invasive fish; *Natrix natrix*; *Natrix tessellata*; semi-aquatic snakes

---

### INTRODUCTION

Grass Snakes (*Natrix natrix*) feed mostly on amphibians (Bruno and Maugeri 1990; Arnold and Ovenden 2002; Gregory and Isaac 2004). However, prey differences among populations, between the sexes, and among age classes within populations have been found in this species (Luiselli et al. 1997, 2005, 2007; Gregory and Isaac 2004; Bilcke et al. 2007). We are not aware of any publication regarding the consumption of allochthonous fish species by *N. natrix*. Another

natricine species, the Dice Snake (*Natrix tessellata*), feeds predominately on fish. As with the Grass Snake, inter-population differences in diet of this species exist. These depend on numerous biotic and abiotic factors such as the differences between males and females and among age (i.e., size) categories (Luiselli et al. 2007). Nevertheless, very few comparative records exist on the consumption of non-native/invasive fishes by this snake (Acipinar et al. 2006).

The variable composition of the prey of natricine snakes implies that these species are quite capable of

surviving in areas where their usual prey decreases dramatically, by shifting to alternative diets, and including non-indigenous species (Gregory and Isaac 2004; Luiselli et al. 2005; Acipinar et al. 2006; King et al. 2006). Inter-population differences in snake diet can be related to climate or habitat characteristics (e.g., Capula and Luiselli 2002; Luiselli et al. 2007). How quickly a shift in diet occurs still remains to be investigated. Diet shifts in the same cohort within a single population, caused by habitat and prey alterations, have seldom been studied.

In general, interactions between non-native prey species and native predators have rarely been addressed (Bulte and Blouin-Demers 2008; Carlsson et al. 2009; Ward-Fear et al. 2009; Li et al. 2011; Robbins et al. 2013). The notable exceptions have been studies regarding the influence of introduced toads to indigenous snake species in Australia (e.g., Shine 2012 and references therein). However, the majority of these studies concern invasive species with chemical “weaponry”; invaders capable of inflicting physical damage are rarely considered.

Analyses and comparisons of Grass and Dice snake diets and their flexibility have not been performed in the Balkans. Our primary aim was to analyze diet composition and its geographic variability (i.e., possible relatedness to the type of habitat) in Grass and Dice snakes. Additional aim of our study was to assess the ratios of prey types (amphibians vs. fish, and native vs. non-native fish species) in the diets of these two snake species under different circumstances.

#### MATERIALS AND METHODS

Grass Snakes occur on three continents: Europe, Asia, and Africa (Arnold and Oviden 2002; Gregory and Isaac 2004). They feed in both aquatic and terrestrial environments. Predominant prey types are amphibians; however, they regularly consume fish (Capula et al. 1994; Filippi et al. 1996; Filippi and Luiselli 2002). Dice Snakes rely almost exclusively on aquatic habitats and their primary prey is fish (Filippi et al. 1996; Bilcke et al. 2007; Capula et al. 2011). Both snakes are common throughout the Palearctic region (Mebert et al. 2011; Shehab et al. 2011).

**Study sites.**—Our first study site, the Bardača Protected Area (Important Bird Area [IBA] and Ramsar site), is a typical marshland. It is located ~50 km north-east of Banja Luka in northern Bosnia and Herzegovina (45°06′ N, 17°26′ E, Fig. 1). The site is at 85–95 m elevation. Its total area before the massive draining that occurred in 2012 was ~3,500 ha. This complex includes fish ponds, floodplain meadows and forests, arable land,



**FIGURE 1.** Localities on the Balkan Peninsula where we studied food preferences of the Grass Snake (*Natrix natrix*) and the Dice Snake (*Natrix tessellata*). In Bosnia and Herzegovina: Bardača Protected Area and Vrbanja River; in Serbia: Pančevački rit wetland; in Montenegro: Skadar Lake; in FYR of Macedonia: Prespa Lake.

and settlements. The ichthyofauna of Bardača comprises 24 species (17% non-native), with two invasive species (Goldfish, *Carassius auratus*, and Brown Bullhead, *Ameiurus nebulosus*) dominating in terms of density and biomass (Vuković et al. 2008; see Appendix). Destructive human activities that have lasted for many decades (Mihajlo Marković, unpubl. report) are continuing. In early 2012, eight of 11 lakes, including our study site, were completely dried out. In the Bardača wetland, we performed the capture-mark-recapture (CMR) surveys of Grass Snakes from March until June in 2011 and 2012 (13 days of fieldwork in total).

Another study site in Bosnia and Herzegovina, in the surroundings of Banja Luka, was the Vrbanja River, a tributary to the Vrbas River (44°47′ N, 17°14′ E, Fig. 1; ~200 m elevation). According to the available literature (Radević 2000; Golub et al. 2012), 21 native fish species are present in this river (see Appendix). Along the river stretch where we conducted our study, strong anthropogenic influences, such as industrial and wastewater discharge and riverbed modifications are evident (Radević 2000; Goran Šukalo, pers. obs.). We performed CMR field studies on Dice Snakes from May to September in 2011 and 2012 (a total of 67 days), along a 2.4 km river stretch. In addition to general population data, we recorded information related to the diet of this species.

The second locality where population (CMR) data on Dice Snakes has been gathered since 2011 is Skadar Lake in Montenegro (40°10′ N, 19°15′ E, Fig. 1; 5 m elevation). This lake, shared between Montenegro and Albania, has the largest surface area in the Balkan Peninsula (391 km<sup>2</sup>), but its average depth is only 6 m. It represents one of the most important centers of biodiversity in southeastern Europe; it has the status of National Park, Ramsar site, and IBA (Radović et al.

2008). There are 47 fish species, mostly cyprinids (see Appendix), of which 14 (29%) are introduced (Cakić and Hristić 1987; Talevski et al. 2009). We made 21 field trips to this site from April to October in 2011 and 2012.

The third locality from which we collected data on the diet of Dice Snakes is Golem Grad Island in Prespa Lake, which, in part, is in the Former Yugoslav Republic of Macedonia (FYROM: 40°52' N, 20°59' E; Fig. 1). This site is ~850 m in elevation and the surface area of the lake is ~250 km<sup>2</sup> with a maximum depth of 55 m (Hollis and Stevenson 1997). The lake and the island are strictly protected within the Galičica National Park. Nevertheless, numerous allochthonous fish species (12 of 23, 52%; see Appendix) have been recorded in the lake (Talevski et al. 2009). Illegal fishing is widespread and common. This particular activity affects Dice Snakes in several ways: it leads to a shortage of food, and many snakes drown entangled in fishing nets set in the shallow waters around the island (Sterijovski et al. 2011). We started a long-term CMR study in 2008 with fieldwork occurring from April to August each year.

The fifth population study on snakes was conducted in the Pančevački rit, an area inundated by the Danube and Tamiš rivers, near Belgrade, Serbia (44°50' N, 20°29' E, Fig. 1; 76 m elevation). This wetland area is under various anthropogenic pressures: amelioration, pollution, construction projects, gravel extraction, burning of dry vegetation, intensive fishing, and introduction of non-native fish species (Cakić and Hristić 1987; Marko Anđelković, pers. obs.). Due to gravel extraction and illegal waste dumping, small water bodies continuously form and disappear; therefore, the composition of the terrain constantly changes. In Pančevački rit, we have studied both species of *Natrix* since 2011; we spent approximately 20 days in the field altogether. In this complex habitat, 35 fish species have been recorded; eight (23%; see Appendix) are introduced (Cakić and Hristić 1987).

**Field procedures.**—At all study sites, we captured snakes by hand and kept them in canvas bags until processing. After measuring and marking, we released them at the exact places of capture. For all individuals we recorded standard measurements: snout-to-vent length (SVL) with a tape measure to 1 mm, and body weight (BM) using an electronic scale, to 1 g. We marked all animals with a standard method of clipping ventral scales (Bonnet et al. 2002; Dorcas and Willson 2009). During this procedure, animals would often regurgitate the ingested prey. Also, we palpated some snakes to make them regurgitate the swallowed food. We used this method only with snakes that had obviously swallowed the prey a short time before capture (i.e., the prey was palpable in the upper portions of the digestive tract); hence, we were sure to obtain an

intact food item. This procedure does not harm the snake (Fauvel et al. 2012).

## RESULTS

We processed between 86 and 5,376 snakes in the five study sites. Sex ratios (M/F) were between 0.28 (Vrbanja River) and 2.77 (Pančevački rit). Percentages of adult animals that we found with prey ranged from 3.51 to 75.00; among juvenile and sub-adult snakes, these percentages varied between 0.63 and 50.00 (Table 1). We calculated the number and percentages of animals with food items in any condition (intact or half-digested, tangible in the stomach and abdomen, or regurgitated) from the total encounter numbers. In all samples, except Grass Snakes from Pančevački rit, there were more females than males that had eaten. This relation was obtained even in populations with male-biased sex ratios. Percentages of sub-adult and juvenile snakes with food were lower than in adults in all samples (Table 1). We obtained identifiable food items from 27 snakes in the wetland complex of Bardača, 19 from the Vrbanja River, eight from the Skadar Lake, and 80 from the Prespa Lake. In the Pančevački rit, we obtained identifiable food items from 18 Grass Snakes and 14 Dice Snakes.

**Diet composition of Grass Snakes.**—In 2011, in the Bardača wetland, we caught the majority of Grass Snakes along the shores of the Necik Lake. Green frogs (*Pelophylax* sp.) dominated (83%) their diet (Table 2). In 2012, Necik Lake was completely drained. Consequently, the snakes disappeared from this primary study area; instead, we searched for them around the adjacent Rakitovac Lake. A small pool formed along one of the Rakitovac Lake drainage canals, and a large number of fish were trapped in it (Fig. 2); the most common species was the Brown Bullhead. The other species present in this trap were the European Perch (*Perca fluviatilis*), the Prussian Carp (*Carassius gibelio*) and several other cyprinids.

In March and April 2012, we captured 70 Grass Snakes along the Rakitovac outlet canal. Of all the snakes we observed feeding or that had recently fed (21), 86% (18 individuals) had consumed Brown Bullhead, and we found three with Prussian Carp (100% of the diet were allochthonous fishes). This was despite the fact that snakes can be seriously injured or even killed by the stiff spines of the dorsal and pectoral fins of the Brown Bullhead (Fig. 3). Another significant observation was that among the 18 Grass Snakes that had caught catfish (*Ameiurus* spp.), 13 were large females (60.2–81.5 cm SVL) and five were males (50.1–58.2 cm SVL). We found one of the smaller females (65 cm SVL) and a small male (50.3 cm) with the spines of catfish

**TABLE 1.** Total numbers and numbers marked (in parentheses) – first three columns, adult sex ratio (M/F), and number and percentages (in parentheses; last three columns) of adult (A), subadult (SA), juvenile (J) Dice Snakes (*Natrix tessellata*) and Grass Snakes (*Natrix natrix*) with intact and half-digested food items (all encounters, captures and recaptures, are considered).

	All Snakes	Adults			Snakes with food		
		F	M	M/F	F	M	SA and J
<i>Natrix tessellata</i>							
Vrbanja River	374 (314)	131 (98)	30 (27)	0.28	20 (15.3%)	3 (10.0%)	31 (14.6%)
Lake Skadar	230 (227)	45 (45)	90 (87)	1.93	10 (22.2%)	16 (17.8%)	13 (13.8%)
Golem Grad Island	5960 (5376)	3369 (2906)	1939 (1826)	0.63	237 (7.03%)	68 (3.51%)	4 (0.63%)
Pančevački rit	162 (161)	14 (13)	36 (36)	2.77	7 (50.0%)	17 (47.2%)	10 (8.93%)
<i>Natrix natrix</i>							
Bardača	115 (108)	57 (53)	46 (43)	0.81	27 (47.4%)	15 (32.6%)	6 (50.0%)
Pančevački rit	86 (86)	20 (20)	12 (12)	0.60	11 (55.0%)	9 (75.0%)	22 (40.7%)

protruding from their bodies, but both snakes were alive and apparently well.

For comparison, Grass Snakes in Pančevački rit (two years pooled; Table 2) consumed mostly anurans: of 14 prey items, 12 (85.7%) were green frogs. One fish species eaten was native to the Balkans and another was the Brown Bullhead, native to eastern North America. In April 2013 we documented two prey species previously unreported for Grass Snakes: the Wall Lizard (*Podarcis muralis*) and the Green Toad (*Pseudepidalea viridis*). In fact, one Grass Snake had ingested both these species.

**Diet composition of Dice Snakes.**—In the four localities where we studied Dice Snake diet, the composition of ichthyofauna differed substantially: from



**FIGURE 2.** A small pool along one of the Rakitovac Lake drainage canals with trapped fish Brown Bullhead, *Ameiurus nebulosus* dominating. (Photographed by Goran Šukalo).

100% autochthonous to 52% allochthonous species (see Appendix), and snakes' prey preferences differed in these circumstances (Table 3). In the Skadar Lake, although the native fish species are dominant (71%), as much as 62% of the diet of Dice Snakes was non-native fishes. In Prespa Lake, the percentages of native and non-native fish species are balanced (48% and 52%, respectively); however, Dice Snakes consumed considerably more autochthonous species (88%). In the Pančevački rit wetland, autochthonous fish species predominate (77%). We recorded 18 fish in the prey of the Dice Snakes from this ecosystem; 17 were European Weatherfish (*Misgurnus fossilis*), an autochthonous species, but we could not identify one species due to the degree of digestion by snake. On two occasions in April 2013, we found Smooth Newts (*Lissotriton vulgaris*) in the prey of Dice Snakes from Pančevački rit. We are not aware of any previous report of newts being consumed by these snakes. In the Vrbanja River, all fish species are autochthonous, and nothing but fish was found in the diet of Dice Snakes, in spite of the presence of huge numbers of green frogs.

## DISCUSSION

Previous studies of European populations of Grass Snakes reported a wide spectrum of prey: amphibians (up to 93%), fish, small reptiles, birds, and rodents (Filippi et al. 1996; Gregory and Isaac 2004; Janev Hutinec and Mebert 2011). Luiselli et al. (2005) found substantial inter-population differences in the composition of the diet of the Grass Snake, dependent on the type of habitat, climate, altitude, and presence of congeneric competitors. The diet is determined by the abundance and availability (i.e., accessibility) of suitable prey (Willson and Hopkins 2011). Non-native species



## Herpetological Conservation and Biology

**TABLE 2.** Grass Snake (*Natrix natrix*) intra-population variation in diet composition due to habitat change in Bardača in two consecutive years and intra-population diet variation in Pančevački rit.

Year	Total No. of prey items	Amphibian prey items	Autochthonous fish prey items	Allochthonous fish prey items
<i>Natrix natrix</i> , Bardača				
2011	6	5 (83%)	0	1 (17%)
2012	21	0%	0	21 (100%)
Pooled 2011 + 2012	27	5 (18.5%)	0	22 (81.5%)
<i>Natrix natrix</i> , Pančevački rit				
2011 spring	3	2 (66.7%)	0	1 (33.3%)
2011 summer	0	0	0	0
2012 spring	10	9 (90%)	1 (10%)	0
2012 summer	1	1 (100%)	0	0
Pooled 2011 +2012	14	12 (85.7%)	1 (7.1%)	1 (7.1%)

of amphibians and fish have been found in the prey of several *Natrix* species (Gregory and Isaac 2004; Santos and García-Cardenete 2005; Alarcos et al. 2009). During our two-year study, we recorded a dramatic change in the prey of the Grass Snake in the Bardača wetland, the habitat that suffered radical alteration. This wetland, although formally protected, is under heavy pressures, and the enormous concentration of alien fish species in its waters could be considered proof of the disturbed balance in this ecosystem (e.g., Cucherousset et al. 2006). After draining the majority of water bodies in Bardača, the snakes started feeding exclusively on allochthonous fishes instead of anurans.

The wide distribution of Grass and Dice snakes (Arnold and Ovenden 2002) illustrates their niche breadth and adaptability under challenging circumstances. Previous records and the findings presented herein demonstrate that snakes can find a way to feed even on fish that can inflict substantial damage or cause death, in some instances (Mills 2002; Santos and

García-Cardenete 2005; Šukalo et al. 2012). One of these harmful fish is the Brown Bullhead, a species of North American origin, which is considered as invasive on other continents (Cakić and Hristić 1987; Declerck et al. 2002). In the native range of ictalurid catfish, there are records of natricine snakes (*Nerodia taxispilota*) that survived massive wounds inflicted by these fish, living for several years, and in some instances, growing considerably (Mills 2002). There are populations within this species in which adults feed almost exclusively on ictalurid catfish (Mills et al. 1995; Mills 2002).

The majority of alien fishes present in the surveyed ecosystems were introduced into Europe and/or the former Yugoslavia 50 to 120 y ago (Cakić and Hristić 1987; Lenhardt et al. 2011; Rutkayová et al. 2013). Therefore, we assume that not all native predators have learned to cope with alien prey and many die trying to eat these fish (Šukalo et al. 2012; data presented herein); however, the potential for overcoming possible perils of feeding on some alien fish is evident. It was experimentally shown that the spines in another ictalurid species (Channel Catfish, *Ictalurus punctatus*) cannot prevent attacks, but they seriously complicate ingestion by gape-limited predators. However, such predators were proven capable of learning to avoid fish with intact spines (Bosher et al. 2006). In the Bardača wetland, the Brown Bullhead was caught mostly by large female Grass Snakes. Among the females that had caught Brown Bullheads, only one was pierced by its spines. This is in accord with the gape size limitations, and with the female-biased sexual dimorphism in body length and relative head size (Luiselli et al. 2007). Thus, it is possible that the snakes that are large enough to swallow a hazardous fish seized an opportunity to feast on the trapped prey. Our data also corroborate previous findings that snakes feed on prey species that are the most abundant, and therefore the easiest to catch under given circumstances (see also King et al. 2006).

Another potentially deadly fish species found in the study areas surveyed during the course of the present



**FIGURE 3.** Grass Snake (*Natrix natrix*) along the Rakitovac outlet canal with half-swallowed Brown Bullhead (*Ameiurus nebulosus*). The fin spine of the fish is protruding from the ventral body wall of the snake. (Photographed by Goran Šukalo).

**TABLE 3.** Ratios of autochthonous to allochthonous fish species in the diet of Dice Snakes (*Natrix tessellata*) from four study sites.

Site	Total No. prey items	Autochthonous fish as prey	Allochthonous fish as prey
Vrbanja River	19	19 (100%)	0 (0%)
Skadar Lake	8	3 (37.5%)	5 (62.5%)
Prespa Lake	80	71 (88.8%)	9 (11.25%)
Pančevački rit	18	18 (100%)	0 (0%)

study is the Pumpkinseed Sunfish (*Lepomis gibbosus*).

This fish also originated in North America and its success in colonizing some freshwater habitats in Europe results from high plasticity in crucial life-history traits (Fox et al. 2007). According to available information (Velikov 2011), there are no published data of Dice Snakes being killed by Pumpkinseed Sunfish. However, its congeners, Grass Snakes and Viperine Snake (*Natrix maura*) can die of the damage inflicted by the sharp fin rays of the Pumpkinseed Sunfish, Brown Bullhead, and other alien fish species (Santos and García-Cardenete 2005; Alarcos et al. 2009; this study). Even without directly damaging the snake, a Pumpkinseed can increase its vulnerability. An experiment with the Southern Watersnake (*Nerodia fasciata*) showed that after ingestion of a fish of inadequate shape (Fig. 4), the mobility of the snake was dramatically reduced (Willson and Hopkins 2011) making it more vulnerable to predators.

In a long-term sense, aquatic snakes can contribute to restoring the previous balance in a disturbed ecosystem, by participating in regulation of the dynamics of invasive fish species populations (Alarcos et al. 2009; Carlsson et al. 2009; Kornis et al. 2012). However, if these fish become the prevalent prey, too many snakes may die struggling to feed. Comparisons of two of our wetland habitats showed that in an ever-changing, polluted, but not drained, wetland (Pančevački rit), Grass Snakes fed mostly on anurans, which is their normal diet. In Bardača, a wetland that had been drained and of which only a small portion remained, the snakes switched to invasive fish. Comparisons of the diets of Dice Snakes showed that in three of four study sites (Vrbanja River, Prespa Lake, and Pančevački rit), Dice Snakes fed predominately on native fish but in Skadar Lake they ate more alien species. Native fish were prevalent in Vrbanja River, Skadar Lake, and Pančevački rit. The discrepancy was found in the Prespa and Skadar lakes. Despite a high proportion of alien fish species, in Prespa the snakes fed mostly on native fish species but in Skadar Lake, despite the prevalence of natives, snakes feed on alien fish species. At present, this result must be considered cautiously, due to the small prey sample size from Skadar Lake. Nevertheless, it can be supposed that in an ancient, deep, and

**FIGURE 4.** Dice Snake (*Natrix tessellata*) starting to swallow a Pumpkinseed Sunfish (*Lepomis gibbosus*) in a small artificial pool near the village Bački Brestovac. (Photographed by Aleksandar Popović).

comparatively well preserved Prespa Lake, the food web is more stable than in other surveyed habitats. This includes Skadar Lake, which is large but shallow, and presumably more prone to variations in abiotic and biotic constituents. Certainly, wider-scope studies are necessary to determine the link between the type and state of a water body, presence of non-native fish species, and relations between various kinds of predators, prey, and their parasites (see Reshetnikov et al. 2013).

Invasions into aquatic ecosystems by various organisms are most likely to take place in disturbed, unstable habitats (Cucherousset et al. 2006, and references therein; Gherardi 2007). Alien fish species can and often do seriously disturb and threaten native ichthyofauna and herpetofauna; their numerous direct and indirect influences have been recorded, but not all are fully understood and explained (Declerck et al. 2002; Kreutzenberger et al. 2008; Copp et al. 2010; Li et al. 2011; Rutkayová et al. 2013). Some snake species have successfully included allochthonous fish and amphibians in their food spectrum (Mullin et al. 2004, Li et al. 2011; Kornis et al. 2012). Further studies of the biology of aquatic snakes would be beneficial in numerous ways. To understand piscivorous snakes, it is necessary to understand native and non-native fish and their biology. Comprehension of the food web and the effects of numerous abiotic and anthropogenic influences on it can lead to the prescription of appropriate preservation and restoration measures for entire aquatic ecosystems (see Matthews et al. 2002; Cucherousset et al. 2006; Phillips and Shine 2006; Carlsson et al. 2009; Shine 2012). Additionally, being the top predators, (semi-)aquatic snakes were long ago recognized as good indicators of the contamination levels of aquatic habitats (Stafford et al. 1976; Ohlendorf et al. 1988; Fontenot et al. 2000). Nevertheless, few studies of contamination of water

bodies have been done with snakes as the main target organism. Future ecological and conservation studies and programs for the restoration of wetlands should consider the feeding habits and behavioral traits of aquatic snakes.

*Acknowledgments.*—Members of the Serbian Herpetological Society “Milutin Radovanović” and biology students and graduates from the Former Yugoslav Republic of Macedonia (FYROM), Montenegro, Croatia, Bosnia and Herzegovina, Serbia, France, and Togo participated in various phases of this research. Special thanks to Dušan Jelić and Ljubomir Pejčić for determining fish species in the field. Prof. Predrag Simonović and Ana Tošić helped with ichthyological literature. Rastko Ajtić provided valuable suggestions during manuscript preparation. Aleksandar Popović kindly allowed us to use his photograph. Our work is permitted by the Ministry of Environment, Mining, and Spatial Planning of Serbia and financed by the Ministry of Education, Science, and Technological Development (permit No. 353-01-46/2012-03, project No. 173043). The authorities of the Ministry of Environment, Mining, and Spatial Planning of FYROM and of the National Park “Galičica” provided permission to work on site (No. 11-4093/5 and 03-246, respectively).

### LITERATURE CITED

- Acipinar, H., O. Gaygusuz, A.S. Tarkan, C. Gursoy, and Z. Al. 2006. Presence of invasive fish species, *Carassius gibelio* (Bloch, 1782) in the diet of the Dice Snake, *Natrix tessellata* (Laurenti, 1768). *Journal of Fisheries and Aquatic Science* 1:213–217.
- Alarcos, G., F. Álvarez-Collado, M.F. Flechoso, J. Madrigal, and M. Lizana. 2009. Peces exóticos de la familia Centrarchidae, un peligro para *Natrix maura*. *Boletín de la Asociación Herpetológica Española* 20:95–97.
- Arnold, E.N., and D.W. Oviden. 2002. *A Field Guide to the Reptiles and Amphibians of Britain and Europe*. 2<sup>nd</sup> Edition. Harper Collins Publishers, London, U.K.
- Bilcke, J., A. Herrel, and P. Aerts. 2007. Effect of prey- and predator size on the capture success of an aquatic snake. *Belgian Journal of Zoology* 137:191–195.
- Bonnet, X., D. Pearson, M. Ladyman, O. Lourdais, and D. Bradshaw. 2002. ‘Heaven’ for serpents? A mark-recapture study of Tiger Snakes (*Notechis scutatus*) on Carnac Island, Western Australia. *Austral Ecology* 27:442–450.
- Bosher, B.T., S.H. Newton, and M.L. Fine. 2006. The spines of the Channel Catfish, *Ictalurus punctatus*, as an anti-predator adaptation: an experimental study. *Ethology* 112:188–195.
- Bruno, S., and S. Maugeri. 1990. *Serpenti d’Italia e d’Europa*. Editoriale Giorgio Mondadori, Milano, Italy.
- Bulte, G., and G. Blouin-Demers. 2008. Northern Map Turtles (*Graptemys geographica*) derive energy from the pelagic pathway through predation on Zebra Mussels (*Dreissena polymorpha*). *Freshwater Biology* 53:497–508.
- Cakić, P., and Dj. Hristić. 1987. The ichthyofauna of Pančevački Rit wetlands (Belgrade) with special reference to the allochthonous fish species. *Bulletin of the Natural History Museum, Belgrade B* 42:103–118.
- Capula, M., L. Rugiero, and L. Luiselli. 1994. Ecological observations on the Sardinian Grass Snake, *Natrix natrix cetti*. *Amphibia-Reptilia* 15:221–227.
- Capula, M., and L. Luiselli. 2002. Feeding strategies of *Elaphe longissima* from contrasting Mediterranean habitats in central Italy. *Italian Journal of Zoology* 69:153–156.
- Capula, M., E. Filippi, L. Rugiero, and L. Luiselli. 2011. Dietary, thermal and reproductive ecology of *Natrix tessellata* in central Italy: a synthesis. *Mertensiella* 18:147–153.
- Carlsson, N.O.L., O. Sarnelle, and D.L. Strayer. 2009. Native predators and exotic prey: an acquired taste? *Frontiers in Ecology and the Environment* 7:525–532.
- Copp, G.H., S. Stakėnas, and J. Cucherousset. 2010. Aliens versus the natives: interactions between introduced Pumpkinseed and indigenous Brown Trout in small streams of southern England. *American Fisheries Society Symposium* 73:347–370.
- Cucherousset, J., J.-M. Paillisson, A. Carpentier, M.-C. Eybert, and J.D. Olden. 2006. Habitat use of an artificial wetland by the invasive catfish *Ameiurus melas*. *Ecology of Freshwater Fish* 15:589–596.
- Declerck, S., G. Louette, T. De Bie, and L. De Meester. 2002. Patterns of diet overlap between populations of non-indigenous and native fishes in shallow ponds. *Journal of Fish Biology* 61:1182–1197.
- Dorcas, M.E., and J.D. Willson. 2009. Innovative methods for studies of snake ecology and conservation. Pp. 5–37 *In* Snakes: Ecology and Conservation. Mullin, S.J., and R.A. Seigel (Eds.). Cornell University Press, Ithaca, New York, USA.
- Fauvel, T., F. Brischoux, M.J. Briand, and X. Bonnet. 2012. Do researchers impact their study populations? Assessing the effect of field procedures in a long term population monitoring of sea kraits. *Amphibia-Reptilia* 33:365–372.
- Filippi, E., M. Capula, L. Luiselli, and U. Agrimi. 1996. The prey spectrum of *Natrix natrix* (Linnaeus, 1758) and *Natrix tessellata* (Laurenti, 1768) in sympatric populations (Squamata: Serpentes: Colubridae). *Herpetozoa* 8:155–164.
- Filippi, E., and L. Luiselli. 2002. Crested Newts *Triturus carnifex* (Laurenti, 1768), form the bulk of the diet in high-altitude Grass Snakes *Natrix natrix* (Linnaeus,

- 1758), of the central Apennines (Caudata: Salamandridae; Squamata: Serpentes). *Herpetozoa* 15:83–85.
- Fontenot, L.W., G.P. Noblet, J.M. Akins, M.D. Stephens, and G.P. Cobb. 2000. Bioaccumulation of polychlorinated biphenyls in ranid frogs and Northern Water Snakes from a hazardous waste site and a contaminated watershed. *Chemosphere* 40:803–809.
- Fox, M.G., A. Vila-Gispert, and G.H. Copp. 2007. Life-history traits of introduced Iberian Pumpkinseed *Lepomis gibbosus* relative to native populations. Can differences explain colonization success? *Journal of Fish Biology* 71:56–69.
- Gherardi, F. (Ed.). 2007. *Biological Invaders in Inland Waters: Profiles, Distribution and Threats*. Invading Nature: Springer Series in Invasion Ecology, Springer, Dordrecht, Netherlands.
- Golub, D., R. Dekić, G. Šukalo, S. Sidak, and S. Lolić. 2012. Ichthyofauna diversity of some Vrbas River tributaries as water quality indicator. Pp. 97–104 *In: The 41th Annual Conference of the Serbian Water Pollution Control Society “WATER 2012” Conference Proceedings*, Divčibare, Serbia. (in Serbian with summary in English).
- Gregory, P.T., and L.A. Isaac. 2004. Food habits of the Grass Snake in southeastern England: is *Natrix natrix* a generalist predator? *Journal of Herpetology* 38:88–95.
- Hollis, G.E., and A.C. Stevenson. 1997. The physical basis of the Lake Mikri Prespa systems: geology, climate, hydrology and water quality. *Hydrobiologia* 351:1–19.
- Janev Hutinec, B., and K. Mebert. 2011. Ecological partitioning between Dice Snakes (*Natrix tessellata*) and Grass Snakes (*Natrix natrix*) in southern Croatia. *Mertensiella* 18:225–233.
- King, R.B., J.M. Ray, and K.M. Stanford. 2006. Gorging on gobies: beneficial effects of alien prey on a threatened vertebrate. *Canadian Journal of Zoology* 84:108–115.
- Kornis, M.S., N. Mercado-Silva, and M.J. Vander Zanden. 2012. Twenty years of invasion: a review of Round Goby *Neogobius melanostomus* biology, spread and ecological implications. *Journal of Fish Biology* 80:235–285.
- Kreutzenberger, K., F. Leprieur, and S. Brosse. 2008. The influence of the invasive Black Bullhead *Ameiurus melas* on the predatory efficiency of Pike *Esox lucius* L. *Journal of Fish Biology* 73:196–205.
- Lenhardt, M., G. Markovic, A. Hegedis, S. Maletin, M. Cirkovic, and Z. Markovic. 2011. Non-native and translocated fish species in Serbia and their impact on the native ichthyofauna. *Reviews in Fish Biology and Fisheries* 21:407–421.
- Li, Y., Z. Ke, S. Wang, G.R. Smith, and X. Liu. 2011. An exotic species is the favorite prey of a native enemy. *PLoS ONE* 6:e24299. doi:10.1371/journal.pone.0024299.
- Luiselli, L., M. Capula, and R. Shine. 1997. Food habits, growth rates, and reproductive biology of Grass Snakes, *Natrix natrix* (Colubridae) in the Italian Alps. *Journal of Zoology*, London 241:371–380.
- Luiselli, L., E. Filippi, and M. Capula. 2005. Geographic variation in diet composition of the Grass Snake (*Natrix natrix*) along the mainland and an island of Italy: the effects of habitat type and interference with potential competitors. *Herpetological Journal* 15:221–230.
- Luiselli, L., D. Capizzi, E. Filippi, C. Anibaldi, L. Rugiero, and M. Capula. 2007. Comparative diets of three populations of an aquatic snake (*Natrix tessellata*, Colubridae) from Mediterranean streams with different hydric regimes. *Copeia* 2007:426–435.
- Matthews, K.R., R.A. Knapp, and K.L. Pope. 2002. Garter Snake distributions in high-elevation aquatic ecosystems: is there a link with declining amphibian populations and non-native trout introductions? *Journal of Herpetology* 36:16–22.
- Mebert, K., A.E. Conelli, M. Nembrini, and B.R. Schmidt. 2011. Monitoring and assessment of the distribution of the Dice Snake in Ticino, Southern Switzerland. *Mertensiella* 18:117–130.
- Mills, M.S., C.J. Hudson, and H.J. Berna. 1995. Spatial ecology and movements of the Brown Water Snake (*Nerodia taxispilota*). *Herpetologica* 51:412–423.
- Mills, M.S. 2002. *Ecology and life history of the Brown Water Snake (Nerodia taxispilota)*. Ph.D. Dissertation, University of Georgia, Athens, Georgia, USA. 231 p.
- Mullin, S.J., H. Imbert, J.M. Fish, E.L. Ervin, and R.N. Fisher. 2004. Snake (Colubridae: *Thamnophis*) predatory responses to chemical cues from native and introduced prey species. *The Southwestern Naturalist* 49:449–456.
- Ohlendorf, H.M., R.L. Hothem, and T.W. Aldrich. 1988. Bioaccumulation of selenium by snakes and frogs in the San Joaquin Valley, California. *Copeia* 1988:704–710.
- Phillips, B.L., and R. Shine. 2006. An invasive species induces rapid adaptive change in a native predator: Cane Toads and Black Snakes in Australia. *Proceedings of the Royal Society* 273:1545–1550.
- Radević, M. 2000. *Ekološki i cenotički odnosi faune riba u srednjem i donjem toku Vrbasa i ribnjaku Bardači*. Monograph, Faculty of Science and Mathematics, Banja Luka, Bosnia and Herzegovina.
- Radović, I., D. Radović, P. Jakšić, G. Džukić, V. Stevanović, Z. Bulić, and V. Bušković. 2008. Skadar Lake region and “target species” species of European conservation concern. *Natura Montenegrina* 7:31–44.
- Reshetnikov, A.N., S.G. Sokolov, I.V. Chikhlyayev, A.I. Fayzulin, A.A. Kirillov, A.E. Kuzovenko, E.N. Protasova, and M.O. Skomorokhov. 2013. Direct and



- indirect interactions between an invasive alien fish (*Percocottus glenii*) and two native semi-aquatic snakes. *Copeia* 2013:103–110.
- Robbins, T.R., N.A. Freidenfelds, and T. Langkilde. 2013. Native predator eats invasive toxic prey: evidence for increased incidence of consumption rather than aversion-learning. *Biological Invasions* 15:407–415.
- Rutkayová, J., R. Biskup, R. Harant, V. Šlechta, and J. Koščo. 2013. *Ameiurus melas* (Black Bullhead): morphological characteristics of new introduced species and its comparison with *Ameiurus nebulosus* (Brown Bullhead). *Reviews in Fish Biology and Fisheries* 23:51–68.
- Santos, X., and L. García-Cardenete. 2005. Introducción de peces en ríos de la Cuenca Mediterránea: una amenaza para sus depredadores. *Boletín de la Asociación Herpetológica Española* 16:50–51.
- Shehab, A.H., A. Al Masri, and Z.S. Amr. 2011. The Dice Snake (*Natrix tessellata*) in Syria: distribution, trade and conservation. *Mertensiella* 18:388–392.
- Shine, R. 2012. Invasive species as drivers of evolutionary change: Cane Toads in tropical Australia. *Evolutionary Applications* 5:107–116.
- Stafford, D.P., F.W. Plapp, and R.R. Fleet. 1976. Snakes as indicators of environmental contamination: relation of detoxifying enzymes and pesticide residues to species occurrence in three aquatic ecosystems. *Archives of Environmental Contamination and Toxicology* 5:15–27.
- Sterijovski, B., R. Ajtić, L. Tomović, S. Djordjević, M. Djurakić, A. Golubović, J. Crnobrnja-Isailović, J.-M. Ballouard, D. Desmont, F. Groupmf, and X. Bonnet. 2011. *Natrix tessellata* on Golem Grad, FYR of Macedonia: a natural fortress shelters a prosperous snake population. *Mertensiella* 18:298–301.
- Šukalo, G., S. Đorđević, D. Dmitrović, and L. Tomović. 2012. Introduced fish *Ameiurus nebulosus* (Le Sueur, 1819): hazard to the Grass Snake *Natrix natrix* (Laurenti, 1768). Photo note. *Hyla Herpetological Bulletin* 2:41–42.
- Talevski, T., D. Milošević, D. Marić, D. Petrović, M. Talevska, and A. Talevska. 2009. Biodiversity of ichthyofauna from Lake Prespa, Lake Ohrid and Lake Skadar. *Biotechnology and Biotechnological Equipment* 23:400–404.
- Velikov, I. 2011. Dice Snake feeds on spiny invasive fish. Photo note. *Mertensiella* 18:447.
- Vuković, D., A. Tursi, R. Carlucci, and R. Dekić. 2008. Ichthyofauna of the wetland ecosystem in the Bardača area (Bosnia and Herzegovina). *Ribarstvo* 66:89–103.
- Ward-Fear, G., G.P. Brown, M.J. Greenlees, and R. Shine. 2009. Maladaptive traits in invasive species: in Australia, Cane Toads are more vulnerable to predatory ants than are native frogs. *Functional Ecology* 23:559–568.
- Willson, J.D., and W.A. Hopkins. 2011. Prey morphology constrains the feeding ecology of an aquatic generalist predator. *Ecology* 92:744–754.



**GORAN ŠUKALO** is a Teaching Assistant at the Faculty of Sciences, University of Banja Luka (Republic of Srpska, Bosnia and Herzegovina) since 2008. He earned a M.Sc. in 2012 for his thesis, Morphological variability and population characteristics of Dice Snake (*Natrix tessellata*) on downflow of Vrbanja river. His research area is herpetology, morphology, and population ecology. (Photographed by Slađana Gvozdenović).



**SONJA ĐORĐEVIĆ** is a Research Associate at the Faculty of Biology, University of Belgrade (Serbia) since 2012. She earned her Ph.D. in 2012 for the thesis, Sexual dimorphism in *Testudo hermanni* from the central part of the Balkan Peninsula. Her research area is herpetology, morphology (and geographic variability in morphological traits), sexual dimorphism and its variability, and population ecology of the Hermann's tortoise (*Testudo hermanni*), vipers, and *Natrix* species in the Balkans. She is a member and one of the founders of the Serbian Herpetological Society "Milutin Radovanović". (Photographed by Oliver Ivanović).



**SLADANA GVOZDENOVIĆ** is a one of founders and a member of NGO Montenegrin Ecologists Society since 2012. She earned a M.Sc. in 2013 for her thesis, Morphological variability and population-ecology characteristics of Dice Snake (*Natrix tessellata*) on Lake Skadar. Her research area is herpetology, morphology, and population ecology. (Photographed by Vuk Iković).



**ALEKSANDAR SIMOVIĆ** is a student at the Faculty of Biology, University of Belgrade. He is focused on population ecology, morphological variability, sexual dimorphism, and conservation biology of amphibians and reptiles. He is a member and one of the founders of the Serbian Herpetological Society "Milutin Radovanović". (Photographed by Milena Krsić).



**MARKO ANĐELKOVIĆ** is a Ph.D. student and a Research Assistant at the Institute for Biological Research "Siniša Stanković", University of Belgrade (Republic of Serbia) since 2013. He graduated in 2011 with the thesis entitled "Comparative analysis of morphological variation and sexual dimorphism in the Dice Snake (*Natrix tessellata*) from two populations from the Former Yugoslav Republic of Macedonia and the Republic of Serbia." His research area is herpetology, morphology, and population ecology. He is a member of the Serbian Herpetological Society "Milutin Radovanović". (Photographed by Ana Golubović).



**VELJKO BLAGOJEVIĆ** is a Ph.D. student at the Institute for Biological Research “Siniša Stanković,” University of Belgrade (Republic of Serbia) since 2013. Veljko graduated in 2013 with his thesis, Catalase protects cardiomyocytes from oxidative damage in diabetes mellitus in rats. His research area is immunomodulation and environmental effects on the murine immune system. (Photographed by Marko Anđelković).



**LJILJANA TOMOVIĆ** is an Associate Professor of Vertebrate Morphology, Systematics, and Phylogeny at the University of Belgrade in the Faculty of Biology. She graduated at the Faculty of Biology, University of Belgrade, where she also completed a Magister and Doctoral thesis on *Vipera ammodytes* systematics and biogeography. She has been studying vipers (*Vipera ammodytes*, *V. berus* and *V. ursinii*) in the central part of the Balkan Peninsula since 1993. Her specialties are herpetology, morphology, systematics, population ecology, and ethology. So far, she has authored or co-authored 32 scientific papers in indexed journals. Since 2005, the main focus of her work has been population studies of *Vipera ursinii* in Macedonia, Montenegro, and Bosnia & Herzegovina, and since 2007, population studies on *Testudo hermanni*, *Natrix tessellata*, *Natrix natrix*, and *Vipera ammodytes* in the central part of the Balkans. Ljiljana is a member and one of the founders of the Serbian Herpetological Society “Milutin Radovanović.” (Photographed by Dragan Arsovski).

**APPENDIX.** Ichthyofauna of the five surveyed localities on the Balkan Peninsula that are potential prey of the Grass Snake (*Natrix natrix*) and the Dice Snake (*Natrix tessellata*). Under each locality: Aut. = autochthonous fish species and All. = allochthonous fish species.

Species	Bardača		Vrbanja		Lake Skadar		Lake Prespa		Pančevački rit	
	Aut.	All.	Aut.	All.	Aut.	All.	Aut.	All.	Aut.	All.
<i>Abramis ballerus</i>										+
<i>Abramis brama</i>	+		+							+
<i>Abramis sapa</i>										+
<i>Acerina cernua</i>	+									+
<i>Acerina schraetser</i>										+
<i>Acipenser naccarii</i>					+					
<i>Acipenser ruthenus</i>										+
<i>Acipenser sturi</i>					+					
<i>Alburnoides bipunctatus</i>			+							
<i>Alburnoides ohridanus</i>					+					
<i>Alburnoides prespensis</i>								+		
<i>Alburnus alburnus</i>	+		+							+
<i>Alburnus belvica</i>								+		

Šukalo et al.—Food Preferences of Two *Natrix* Species.

APPENDIX. Ichthyofauna of the five surveyed localities on the Balkan Peninsula that are potential prey of the Grass Snake (*Natrix natrix*) and the Dice Snake (*Natrix tessellata*). Under each locality: Aut. = autochthonous fish species and All. = allochthonous fish species.

Species	Bardača		Vrbanja		Lake Skadar		Lake Prespa		Pančevački rit	
	Aut.	All.	Aut.	All.	Aut.	All.	Aut.	All.	Aut.	All.
<i>Alburnus scoranza</i>					+					
<i>Alosa falax</i>					+					
<i>Ameiurus nebulosus</i>							+			+
<i>Anguilla anguilla</i>					+		+			
<i>Aspius aspius</i>									+	
<i>Barbatula zetensis</i>					+					
<i>Barbus balcanicus</i>			+							
<i>Barbus barbus</i>			+							
<i>Barbus prespensis</i>							+			
<i>Barbus rebeli</i>					+					
<i>Blicca bjoerkna</i>	+								+	
<i>Carassius carassius</i>									+	
<i>Carassius gibelio</i>		+					+	+		+
<i>Chalcalburnus chalcoides</i>	+								+	
<i>Chondrostoma nasus</i>	+		+						+	
<i>Chondrostoma prespense</i>							+			
<i>Chondrostoma scodrensis</i>					+					
<i>Citharus linguatulus</i>					+					
<i>Cobitis elongatoides</i>		+								
<i>Cobitis ohridana</i>				+						
<i>Cobitis meridionalis</i>						+				
<i>Cobitis taenia</i>	+						+			
<i>Cottus gobio</i>		+								
<i>Ctenopharyngodon idella</i>					+		+		+	
<i>Cyprinus carpio</i>	+	+	+		+		+			
<i>Dicentrarchus labrax</i>				+						
<i>Esox lucius</i>	+						+			
<i>Gambusia holbrooki</i>					+		+			
<i>Gasterosteus gymnurus</i>				+						
<i>Gobio gobio</i>		+								
<i>Gobio obtusirostris</i>	+	+								
<i>Gobio skadrensis</i>				+						
<i>Hucho hucho</i>		+								

## Herpetological Conservation and Biology

**APPENDIX.** Ichthyofauna of the five surveyed localities on the Balkan Peninsula that are potential prey of the Grass Snake (*Natrix natrix*) and the Dice Snake (*Natrix tessellata*). Under each locality: Aut. = autochthonous fish species and All. = allochthonous fish species.

Species	Bardača		Vrbanja		Lake Skadar		Lake Prespa		Pančevački rit	
	Aut.	All.	Aut.	All.	Aut.	All.	Aut.	All.	Aut.	All.
<i>Hypophthalmichthys molitrix</i>					+		+			+
<i>Hypophthalmichthys nobilis</i>					+					+
<i>Ictalurus nebulosus</i>	+				+					+
<i>Lampetra fluviatilis</i>								+		
<i>Lepomis gibbosus</i>	+						+			+
<i>Leuciscus idus</i>		+						+		
<i>Leuciscus leuciscus</i>	+									
<i>Liza ramada</i>				+						
<i>Megalobrama terminalis</i>						+				
<i>Misgurnus fossilis</i>	+							+		
<i>Mugil cephalus</i>				+						
<i>Mylopharyngodon piceus</i>						+				
<i>Oncorhynchus mykiss</i>						+		+		
<i>Pachychilon pictum</i>				+						
<i>Parabramis pekinensis</i>								+		
<i>Pelagus minutus</i>				+						
<i>Pelagus prespensis</i>						+				
<i>Perca fluviatilis</i>	+			+				+		
<i>Pleuronectes flessus</i>				+						
<i>Phoxinus phoxinus</i>		+								
<i>Pleuronectes flessus</i>				+						
<i>Pomatoschistus montenegrensis</i>				+						
<i>Pseudorasbora parva</i>		+		+		+		+		
<i>Rhodeus amarus</i>	+		+	+		+				
<i>Rhodeus sericeus</i>								+		
<i>Romanogobio kesslerii</i>		+								
<i>Rutilus ohridanus</i>				+						
<i>Rutilus prespensis</i>						+				
<i>Rutilus pigus</i>		+								
<i>Rutilus rutilus</i>	+							+		
<i>Salaria fluviatilis</i>				+						
<i>Salmo farioides</i>				+						



Šukalo et al.—Food Preferences of Two *Natrix* Species.

APPENDIX. Ichthyofauna of the five surveyed localities on the Balkan Peninsula that are potential prey of the Grass Snake (*Natrix natrix*) and the Dice Snake (*Natrix tessellata*). Under each locality: Aut. = autochthonous fish species and All. = allochthonous fish species.

Species	Bardača		Vrbanja		Lake Skadar		Lake Prespa		Pančevački rit	
	Aut.	All.	Aut.	All.	Aut.	All.	Aut.	All.	Aut.	All.
<i>Salmo letnica</i>					+					
<i>Salmo marmoratus</i>			+							
<i>Salmo peristericus</i>						+				
<i>Salmo trutta</i> , m. <i>fario</i>	+									
<i>Salmothymus zetensis</i>			+							
<i>Salvelinus fontinalis</i>				+						
<i>Scardinius erythrophthalmus</i>	+							+		
<i>Scardinius knezevici</i>			+							
<i>Silurus glanis</i>	+					+	+			
<i>Squalius cephalus</i>	+	+								
<i>Squalius prespensis</i>						+				
<i>Squalius squalus</i>			+							
<i>Stizostedion lucioperca</i>	+							+		
<i>Telestes montenegrinus</i>			+							
<i>Thymallus thymalus</i>		+		+						
<i>Tinca tinca</i>	+			+		+	+			
<i>Umbra krameri</i>								+		
<i>Vimba vimba</i>		+						+		
Total Number	20	4	21	0	34	14	11	12	27	8
Percentage Auto. and Allo.	83	17	100	0	71	29	48	52	77	23